


# Evolution in Sweden 2025

January 13 – 15, 2025, Linköping



Dear conference participant,

We look forward to welcoming you to the Evolution in Sweden conference in Linköping, 13<sup>th</sup>-15<sup>th</sup> January, 2025! This pdf contains the abstracts (oral and poster) for the conference, a preliminary (hopefully final, but any changes will be updated on a google docs excel file and sent to everyone) program.

The conference will be held in the C-building at the Valla (main) campus of Linköping University. The conference will be held in the lecture theatre C4, with the posters set-up in the foyer-area outside C4. Registration is from 9.30 until 13.00 outside the lecture hall on the 13<sup>th</sup> January.

There is a lunch venue booked at the University restaurant Kårallen (see map on the next page), which is situated in the next building down from the C-building (two minutes walk from the lecture hall). We have booked a reservation for all those who indicated that they would like a reservation (and can most likely accommodate any who had forgotten to book but changed their minds). The cost of lunch is at everyone's own expense. In addition, we will provide tea, coffee and biscuits at the fika breaks, with this generously provided by Qiagen (stop by their booth at the conference and say hello!). We would also like to thank the SciLifeLab for generously hosting the plenary speakers and for hosting a DDLS speaker session on the 14<sup>th</sup>!

We have a poster session on the 13<sup>th</sup> and 14<sup>th</sup> outside the lecture theatre, and have also highlighted a conference pub in town (O'Leary's). This was picked because of its size, but we don't have any actual reservations (but it is pretty huge on the inside). It is located in the main square (Storatorget), and several other pubs are also located on the main square or on the next street (Ågatan) for those who wise to meet for a drink in the evenings.

We hope you all have a lovely Christmas and New Year, and look forward to all the exciting talks and seeing everyone in the new year.


Best

The Evolution in Sweden Linköping organizing committee



Dominic Wright (chair), Rie Henriksen, Per Jensen, Urban Friberg and Krzysztof Bartoszek

### Getting to the venue:

From the bus station (next to the train station), you can take the bus numbers 4, 540, 39, otherwise it is around a one hour walk



### Maps showing the conference location (C building), venue location (C4 lecture hall), nearest bus stop, and lunch venue



## EVOLUTION IN SWEDEN LINKÖPING 2025 FULL PROGRAM

### January

13th

**09:30 On-site registration outside C4**

**12:00 Lunch**

12:45 Welcome

**PLENARY - Tom**

**13:00 Gilbert**

14:00	Jessica Abbott	Change in the G-matrix as a result of female-limited X chromosome evolution
14:15	Ted Morrow	The adaptive value of recombination in resolving intralocus sexual conflict by gene duplication
14:30	Ingo Müller Mafalda Sousa	Hybridisation patterns in the birds-of-paradise genus <i>Paradisaea</i>
14:45	Ferreira	The role of introgression for adaptation of Atlantic herring to the Baltic sea

**15:00 FIKA**

**Sponsored by Qiagen**

15:30	Krzysztof Bartoszek	Fast inference for multivariate phylogenetic comparative methods
15:45	Ivan Prates	Unraveling the genomic underpinnings of parallel evolution in Mediterranean wall lizards
16:00	Anna Qvarnström	Inheritance of material wealth in a natural population
16:15	Enya Van Poucke	Domestication and Early Stress Resilience: Insights from White Leghorns and Red Junglefowl.
16:30	Alexandre Rego	Selective hearts, selected genes: genomic consequences of different mating regimes
16:45	Anna Runemark	The genomic basis of adaptation to a new host plant
17:00	Lars Rönnegård	Could selection for decreased variability in milk yield lead to asocial cows?
17:15	Pamela Santana	Evolution of flower shape in a coevolved system
17:30	Hanna Sigeman	A novel supergene explains alternative reproductive strategies in the ant <i>M. ruginodis</i>

**17:45 POSTER SESSION**

**20:00 Conference Pub O'Leary's Storaorget**

### January 14th

**09:00 PLENARY - Kees Van Oers**

**10:00 FIKA**

**Sponsored by Qiagen**

**DDL Session**

10:30	Jacob Höglund	The SciLifeLab Planetary Biology Capability
10:45	Lisandro Milocco	Toward predictive evo-devo: a data-driven approach to modeling phenotypic variation
11:00	Nikolay Oskolkov	aMeta: a computational method for data-driven ancient metagenomic analysis
11:15	Masahito Tsuboi	Dialectical relationships between microevolution and macroevolution
11:30	José Cerca	Step by step, an adaptive radiation is made: Empirical evidence for the 'least action effect'

11:45	Lennart Winkler	How ecologically relevant resource distributions influence condition-dependent trade-offs
<b>12:00</b>	<b>LUNCH</b>	
13:15	Staffan Bensch	Identifying the elusive migration genes
13:30	David Wheatcroft	TBC
13:45	David Berger	Strong Selection, but low repeatability: Temperature-specific effects on genomic predictions of adaptation
14:00	Folmer Bokma	Contrasting evolution in incipient versus established species
14:15	Diego Stingo Hirmas	Prenatal maternal stress underlies multi-hierarchical behavioural plasticity
14:30	Marianne Dehasque	From Wild to Tamed and Back: the evolutionary history of Sardinian sheep
14:45	Eran Elhaik	Why All PCAs in Evolutionary Biology are Likely Wrong
<b>15:00</b>	<b>FIKA</b>	<b>Sponsored by Qiagen</b>
15:30	Turid Everitt	Investigating the role of meiotic recombination in social insects - Termites adding an important piece to the puzzle
15:45	John Fitzpatrick	The evolution of conspicuous markings in skates and rays
16:00	Samyuktha Rajan	Mechanisms of mixed singing and species recognition in sympatric flycatchers
16:15	Ana Gomes	Experimental examination of the neural background to cognitive variation across species in Poeciliid fishes
16:30	Jake Goodall	Atlantic Herring - Population trends and perspectives from the Baltic Sea
16:45	Chloé Haberkorn Christina Hansen	Adaptive benefits of hybridization in <i>Saccharomyces</i> yeast under thermal stress: genomic and fitness analyses
17:00	Wheat	The importance of standing ancestral variation in behavioural evolution
17:15	Joshua Hufton	Epigenetic regulation of immune response in the Swedish sand lizard ( <i>Lacerta agilis</i> )
<b>17:30</b>	<b>POSTER SESSION</b>	
<b>20:00</b>	<b>Conference Pub</b>	<b>O'Leary's Storatorget</b>

**January  
15th**

<b>09:00</b>	<b>PLENARY - Susan Johnston</b>	
<b>10:00</b>	<b>FIKA</b>	<b>Sponsored by Qiagen</b>
10:30	Zeeshan Syed	A study of the aging transcriptome from an adaptive perspective
10:45	Julia Höglund	Characterisation and quantification of deleterious genetic variants in non-model organisms: from present to extinct species
11:00	Martin Johnsson	Loss of genetic variation and establishment of damaging variants in effectively small populations
11:15	Cheng Ma	The adaptive evolution of herring to different salinities
11:30	R. Axel W. Wiberg	Testing genomic predictions of sexually antagonistic coevolution: Natural variation in the sex-peptide network in <i>Drosophila melanogaster</i>
11:45	Dick Moberg	Speciation and Temporal Dynamics of Female Sperm Management
<b>12:00</b>	<b>LUNCH</b>	

**Abstract number: Plenary****Title:** Domestication hologenomics – how did our ancestors tame the wolf?**Name:** Prof. Tom Gilbert**University:**

Comparisons of the genomes of contemporary domestic animals and plants with those of their wild relatives, have provided a wealth of insights into not only when and where our ancestors started the process, but also what specific genetic variants are key to modern phenotypes. Furthermore, once coupled to palaeogenomic data, such datasets can also even reveal the order in which such variants arose, shedding further insights into the process itself. However, while there is no doubt that we have learnt much about domestication in general, and indeed for most domestic species we can clearly document the genetic basis of why the end product differs from the start, I argue that there may be certain processes that were involved that have been largely overlooked, in particular related to the so-called hologenome.

**Biography:**

Tom Gilbert is Professor of Palaeogenomics at the University of Copenhagen, Professor II at NTNU University Museum, and Director of the DNRF Center for Evolutionary Hologenomics. He holds a PhD in the study of ancient DNA from the University of Oxford, and has been active in both trying to develop methods to both expand the potential of ancient DNA to our understanding of the past, as well as more recently leading research aimed at revisiting our understanding of ecological and evolutionary processes using hologenomic techniques – ie the integrative approach of combining host genomes with those of their microbiomes.



**Abstract number: Plenary**

**Title:** The epigenetics of Animal Personality

**Name:** Prof. Kees Van Oers

**University:** Netherlands Institute of Ecology/ Wageningen University

**Abstract:**

Epigenetic mechanisms are those molecular mechanisms that affect gene expression without changing the DNA sequence. The value of epigenetic mechanisms is increasingly recognized, also in relation to questions in ecology and evolution. However, epigenetic research related to behavioural variation in the natural habitat is still in its infancy. The flexible nature of epigenetic marks opens the possibility that such changes are adaptive, while at the same time may simply be the consequence of environmental variation. Hence, changes in epigenetic marks can function as switches in order to help an organism develop, as signals of aging via accumulation of methylation over time, but they also may aid organisms to cope with changing circumstances throughout their lifetime. In this presentation, I use data of our great tit system to show examples of how tissue and cell-specific epigenetic patterns, mainly focusing on DNA methylation, may affect behavioral phenotypes such as exploratory behavior and reversal learning performance. I discuss what the role is of epigenetic mechanisms for behavioral adaptation to changing environments.

**Biography:**

Kees van Oers is an evolutionary behavioural ecologist. He currently holds a chair in Animal Personality at Wageningen University and is senior scientist at the Netherlands Institute of Ecology (NIOO-KNAW). His research focuses on explaining the causes and consequences of individual variation in animal behaviour, mainly cognitive and animal personality traits. He does this from an ecological and evolutionary point of view, at the crossroads between behavioural ecology and behaviour genetics. His goal is to find answers to fundamental and strategic questions related to individual responses to changing environments. This is relevant since the world and therefore the environment is constantly changing partly due to human influences affecting biodiversity at all functional levels.



**Abstract number: Plenary**

**Title:** The causes and consequences of sex differences in recombination rates.

**Name:** Dr. Susan Johnston

**University:** Edinburgh University

**Abstract:**

The rate of meiotic recombination often shows large differences between the sexes. It can be strongly female-biased (humans), strongly male-biased (macaques/sheep) or somewhere in between. Understanding why this happens is key to understanding the evolution of recombination rates, yet the causes and consequences of this phenomenon remain unknown. This talk will focus on our most recent work in house sparrows (*Passer domesticus*), with broader context from work in mammals and fish. We use genomic data in large pedigrees to characterise individual recombination rates and landscapes to: (a) investigate the heritability and genomic basis of variation in recombination rates; (b) identify genomic correlates with fine-scale sex-differences in recombination landscapes; and (c) use genomic prediction approaches to understand the relationship between recombination and fitness within each sex. Our work provides a foundation for broader understanding of the vast diversity of recombination rates in eukaryotic genomes.

**Biography:**

Susan Johnston is an evolutionary quantitative geneticist based at the University of Edinburgh, Scotland, where she is a Senior Lecturer and Royal Society University Research Fellow. Her research centres on using genomic information to understand selection and evolution in both wild and domesticated populations. At the moment, her group focusses on questions related to genomic signatures of sexual conflict, the maintenance of genetic variation in immunity, and the causes and consequence of recombination landscape variation. She does this by taking advantage of long-term ecological datasets and deep pedigrees with genomic data in mammals, birds and fish.





**Abstract number:** O1

**Title:** Change in the G-matrix as a result of female-limited X chromosome evolution

**Name:** Jessica Abbott

**University:** Lund University

**Abstract:**

Genetic variation on the X chromosome is likely to be subject to unequal sex-specific selection pressures. This is due to its unequal transmission pattern between the sexes, combined with the potential for stronger sexual selection in one sex. Nevertheless, the evolution of sexual dimorphism may be constrained by the shared genetic basis of homologous traits. After more than 147 generations of female-limited X chromosome (FLX) experimental evolution we estimated the phenotypic response to selection and quantitative genetic parameters for three homologous traits in *Drosophila melanogaster* (wing length, desiccation resistance, and locomotory activity). We expected that changes in allele frequencies associated with phenotypic feminization as a result of FLX evolution might modify the underlying genetic basis of these traits. We found evidence of a significant phenotypic change in desiccation resistance in the FLX selection regime, but contrary to our expectations, this change was not consistent with feminization. We also found some evidence of consistent changes in the genetic architecture of locomotory activity in the FLX selection regime, which was surprising given that this trait did not show any evidence of a phenotypic change. We outline a number of possible explanations for this result.

**Authors:** Manat, Yesbol, Lund University, Sweden/Houston University, USA  
Tarka, Maja, Lund University, Sweden  
Abbott, Jessica, Lund University, Sweden

**Key words:** quantitative genetics, sexual conflict, experimental evolution, sex chromosomes

**Email:** [jessica.abbott@biol.lu.se](mailto:jessica.abbott@biol.lu.se)

**Abstract number:** O2

**Title:** The adaptive value of recombination in resolving intralocus sexual conflict by gene duplication

**Name:** Ted Morrow

**University:** Karlstad University

**Abstract:**

We investigate whether recombination can play a role in resolving intralocus sexual conflict. Errors during recombination resulting in gene duplications may provide a long-term evolutionary advantage if those loci also experience sexually antagonistic (SA) selection since, after duplication, sex-specific expression profiles will be free to evolve, thereby reducing the load on population fitness. Using data on candidate SA loci from *Drosophila melanogaster* and humans we found single-nucleotide polymorphisms (SNPs) with net-positive effects across the two sexes occurred at higher frequencies than SNPs with net-negative effects. In flies, higher recombination rates were associated with more intense levels of sexual conflict and genes with paralogs occur in regions with higher recombination rates. Genes experiencing higher levels of conflict also showed both a higher proportion and higher numbers of paralogs. Our results support a possible route towards resolution of an adaptive conflict via gene duplication that is facilitated by higher recombination rates.

**Authors:** Jon A Harper, University of Sussex, UK Ted Morrow, Karlstad University, Sweden

**Key words:** Sexual conflict, recombination, resolution, gene duplication

**Email:** ted.morrow@kau.se

**Abstract number:** O3

**Title:** Hybridisation patterns in the birds-of-paradise genus *Paradisaea*

**Name:** Ingo Müller

**University:** Swedish Museum of Natural History / Stockholm University

**Abstract:**

Hybridisation offers a great opportunity to study speciation, by allowing us to investigate mechanisms driving reproductive isolation between populations. Sexual selection is one mechanism that could ultimately reduce the prevalence of hybridization and promote speciation. However, interestingly, our previous research has shown that hybridization may be surprisingly common in taxa with extreme forms of sexual selection, such as lekking Birds-of-Paradise. In this study, we investigated hybridisation patterns within the genus *Paradisaea* using a whole-genome dataset of more than 100 samples, primarily obtained from historical samples (hDNA). Through comprehensive population genetic and phylogenetic analyses, we assessed the prevalence of hybridisation within the genus and found significant mixing between species with sympatric distributions on New Guinea's mainland. We also identified patterns of introgression driven by backcrossing deep into the parental species' distributions. Finally, our study further highlights the importance of museum collections to study genomic change in species that are difficult to sample in the wild.

**Authors:** Müller, Ingo A. (Department of Bioinformatics and Genetics, Swedish Museum of Natural History, Stockholm, Sweden / Division of Systematics and Evolution, Department of Zoology Stockholm University) Waqar, Abdul (Department of Bioinformatics and Genetics, Swedish Museum of Natural History, Stockholm, Sweden) Thörn, Filip (Department of Bioinformatics and Genetics, Swedish Museum of Natural History, Stockholm, Sweden) Jønsson, Knud Andreas (Department of Bioinformatics and Genetics, Swedish Museum of Natural History, Stockholm, Sweden) Irestedt, Martin (Department of Bioinformatics and Genetics, Swedish Museum of Natural History, Stockholm, Sweden) Blom, Mozes P.K. (Museum für Naturkunde, Leibniz Institut für Evolutions- und Biodiversitätsforschung, Berlin, Germany)

**Key words:** Museomics, Hybridisation, Population Genomics, Phylogenetics, Bioinformatics

**Email:** Ingo.Mueller94@gmail.com

**Abstract number:** O4

**Title:** aMeta: a computational method for data-driven ancient metagenomic analysis

**Name:** Nikolay Oskolkov

**University:** Lund University

**Abstract:**

Analysis of microbial data from archaeological samples is a growing field with great potential for understanding ancient environments, lifestyles, and diseases. However, high error rates have been a challenge in ancient metagenomics, and the availability of computational frameworks that meet the demands of the field is limited. Here, we propose aMeta [1], an accurate metagenomic profiling workflow for ancient DNA which is designed to minimize the amount of false discoveries and computer memory requirements. Using simulated data, we benchmark aMeta against a current state-of-the-art workflow and demonstrate its superiority in microbial detection and authentication, as well as substantially lower usage of computer memory.

[1] Zoé Pochon, Nora Bergfeldt, Emrah Kırdök, Mário Vicente, Thijessen Naidoo, Tom van der Valk, N. Ezgi Altınışık, Maja Krzewińska, Love Dalen, Anders Götherström, Claudio Mirabello, Per Unneberg and Nikolay Oskolkov, aMeta: an accurate and memory-efficient ancient Metagenomic profiling workflow, *Genome Biology* 2023, 24 (242), <https://doi.org/10.1186/s13059-023-03083-9>

**Authors:** Nikolay Oskolkov, Biology Department, Science for Life Laboratory, National Bioinformatics Infrastructure Sweden, Lund University, Lund, Sweden

**Key words:** ancient DNA, metagenomics, microbiome, ancient metagenomics, microbial pathogens

**Email:** Nikolay.Oskolkov@biol.lu.se

**Abstract number:** O5

**Title:** Fast inference for multivariate phylogenetic comparative methods

**Name:** Krzysztof Bartoszek

**University:** Linköping University

**Abstract:**

Co-adaptation is key to understanding species evolution. Different traits have to function together so that the organism can work as a whole. Hence, all changes to environmental pressures have to be coordinated. Since 2019, we have been developing R packages that are able to very efficiently do estimation for general, multivariate Gaussian processes realized over a phylogenetic tree. Using the likelihood computational engine PCMBase (on CRAN) mvSLOUCH (on CRAN) now is able to efficiently work with multivariate phylogenetic Ornstein-Uhlenbeck models. The user is able to propose specific model parameterizations that correspond to particular hypotheses about relationships between traits. In the talk I will discuss how one can setup different hypotheses concerning relationships between the traits in terms of model parameters and discuss particular aspects of the estimation procedures, based on our recent wide simulation studies. The software's possibilities will be illustrated by considering the evolution of functional traits in a dataset of 1252 angiosperms.

**Authors:** Krzysztof Bartoszek

**Key words:** phylogenetic comparative methods, R, maximum likelihood inference, pruning algorithm, angiosperms

**Email:** krzysztof.bartoszek@liu.se

**Abstract number:** O6

**Title:** Unraveling the genomic underpinnings of parallel evolution in Mediterranean wall lizards

**Name:** Ivan Prates

**University:** Lund University

**Abstract:**

Shared phenotypes among distantly related species provide compelling evidence of parallel evolution in response to common ecological pressures or developmental biases. Nevertheless, the genetic mechanisms underlying parallel evolution remain poorly known. Here, we perform a comparative phenomic and genomic study of Mediterranean wall lizards (genus *Podarcis*). We show that several *Podarcis* species independently evolved a similar suite of consistently correlated characters, including coloration, morphology, and behavior. In each species, this ‘syndrome’ is present in only some populations, which enabled us to investigate the genomic basis of the repeated evolution of complex phenotypes. To this end, we employed whole-genome re-sequencing and genome scans to identify and compare loci associated with the syndrome across species, finding both shared and unique outlier loci. This combined approach provides insights into the factors underpinning repeated evolutionary outcomes – a pervasive feature of life on Earth.

**Authors:** Prates Ivan, Lund University, Sweden Abalos Javier, Lund University, Sweden Uller Tobias, Lund University, Sweden Feiner Nathalie, Lund University, Sweden

**Key words:** Parallel evolution, phenotype, coloration, genomics, adaptation

**Email:** [ivanprates@gmail.com](mailto:ivanprates@gmail.com)

**Abstract number:** O7

**Title:** Inheritance of material wealth in a natural population

**Name:** Anna Qvarnström

**University:** Uppsala University

**Abstract:**

Evolutionary adaptation occurs when individuals vary in access to fitness-relevant resources and these differences in “material wealth” are heritable. It is typically assumed that the evolutionary outcomes only depend on heritable variation in the phenotypic abilities needed to acquire material wealth. Here, we scrutinize this assumption by investigating additional mechanism underlying inheritance of material wealth in collared flycatchers. A genome-wide-association analyses reveals a high genomic heritability ( $h^2=0.405\pm 0.08$ ) of access to caterpillar larvae, a fitness-relevant resource, in the birds’ breeding territories. Analyses of the top 100 SNPs associated with this material wealth reveal little evidence for corresponding associations with causal phenotypic acquisition abilities. Simulations, experimental and long-term monitoring data instead imply that heritability of material wealth is largely explained by philopatry that causes within-population genetic structure across a heterogeneous landscape. Therefore, allelic variants associated with high wealth may spread in the population without having causal connections to traits promoting local adaptation.

**Authors:** Murielle Ålund<sup>1</sup>, *S. Eryn McFarlane*<sup>2</sup>, *Arild Husby*<sup>3</sup>, *Jonas Knape*<sup>4</sup>, *Tomas Pärt*<sup>4</sup>, *Päivi Sirkkiä*<sup>5</sup>, *Franz J. Weissing*<sup>6</sup>, *David Wheatcroft*<sup>7</sup>, *Yishu Zhu*<sup>1</sup> and *Anna Qvarnström*<sup>1</sup>

Affiliation and e-mail address <sup>1</sup>Department of Ecology and Genetics (IEG), Division of Animal Ecology, Uppsala University, Norbyvägen 18D, SE-752 36 Uppsala, Sweden Email: Murielle.Alund@ebc.uu.se Email: Yishu.Zhu@ebc.uu.se Email: Anna.Qvarnstrom@ebc.uu.se

<sup>2</sup>Department of Biology, York University, Toronto, Ontario, Canada Email: emcfar@yorku.ca

<sup>3</sup>Department of Ecology and Genetics (IEG), Division of Evolutionary Biology, Uppsala University, Norbyvägen 18D, SE-752 36 Uppsala, Sweden Email: arild.husby@ebc.uu.se

<sup>4</sup>Department of Ecology, Swedish University of Agricultural Sciences, SLU, Box 7044, SE-75007 Uppsala, Sweden Email: Jonas.Knape@slu.se Email: Tomas.Part@slu.se

<sup>5</sup>Finnish Environmental Institute, Latokartanonkaari 11, 00790 Helsinki, Finland Email: Paivi.Sirkia@syke.fi

<sup>6</sup>Groningen Institute for Evolutionary Life Sciences, University of Groningen, Nijenborgh 7, 9747 AG Groningen, The Netherlands Email: f.j.weissing@rug.nl

7Department of Zoology, Stockholm University, Svante Arrhenius väg, SE-106  
91 Stockholm, Sweden Email: David.Wheatcroft@zoologi.su.se

**Key words:** site fidelity, social inheritance, landscape genomics, heritability,  
GWAS, animal breeding model,

**Email:** anna.qvarnstrom@ebc.uu.se



**Abstract number:** O8

**Title:** Mechanisms of mixed singing and species recognition in sympatric flycatchers

**Name:** Samyuktha Rajan

**University:** Stockholm University

**Abstract:**

Assortative mating relies heavily on recognizing conspecifics, a process often mediated by acoustic signals in birds. On the island of Öland, where pied and collared flycatchers occur in sympatry, pied flycatchers deviate from their species-specific songs by producing a converged ‘mixed’ song that partly resembles collared flycatcher song. This asymmetric song convergence has previously been shown to increase interspecies mating errors. However, it remains unclear whether pied flycatchers incorporate collared flycatcher syllables into their songs (true mixed singing) or simply modify their own songs to approximate collared flycatchers’ due to broad learning predispositions. To disentangle these hypotheses, we conducted an in-depth acoustic analysis to trace the origins of syllables within mixed songs. This study provides insights into the mechanisms underlying asymmetric song learning in natural populations, which can ultimately help us predict how song learning may impact reproductive isolation and the evolutionary trajectory of these species.

**Authors:** Rajan Samyuktha, Stockholm University, Sweden Segami Carolina, Uppsala University, Sweden and Duke University, United States of America Ålund Murielle, Uppsala University, Sweden Qvarnström Anna, Uppsala University, Sweden Wheatcroft David, Stockholm University, Sweden

**Key words:** Speciation, cultural evolution, song learning, *Ficedula*

**Email:** samyuktha.rajan@zoologi.su.se

**Abstract number:** O9

**Title:** Selective hearts, selected genes: genomic consequences of different mating regimes

**Name:** Alexandre Rego

**University:** Uppsala University

**Abstract:**

Sexual selection is theorized to play a pivotal role in shaping genomic variation and architecture, either by purging deleterious alleles or by maintaining genetic diversity through balancing selection across the sexes. Despite its theoretical importance, empirical tests of this hypothesis remain limited. Here, we conducted experimental evolution spanning 90 generations in seed beetles maintained under three alternative mating regimes: polygamy (intense sexual selection), monogamy (eliminating sexual selection), and male-limited selection (restricting selection to males). By comparing these regimes, we investigated how sexual selection interacts with natural selection to shape genomic variation, focusing on estimating the relative importance of purging of deleterious alleles and the maintenance of genetic variation through sexually antagonistic selection. By integrating transcriptomic and genomic analyses, we provide a comprehensive view of how mating systems influence genomic variation and pinpoint genetic mechanisms underpinning sex-specific adaptation and conflict.

**Authors:** Alexandre Rego, Uppsala University, Sweden Mareike Koppik, Martin-Luther University Halle-Wittenberg, Germany Julian Baur, Uppsala University, Sweden David Berger, Uppsala University, Sweden

**Key words:** sexual selection, experimental evolution, genetic variation, genomic architecture, *Callosobruchus maculatus*

**Email:** [denovorego@gmail.com](mailto:denovorego@gmail.com)

**Abstract number:** O10

**Title:** The genomic basis of adaptation to a new host plant

**Name:** Anna Runemark

**University:** Lund University

**Abstract:**

While populations are known to rapidly adapt to new ecological niches, how divergent ecological adaptation translates into reproductive isolation and ultimately speciation is a topic of current debate. Here, we use *Tephritis conura* peacock flies adapted to different thistle host plants to uncover the genomic basis enabling the use of different thistle host plants, also in sympatry. We document a large inversion on chromosome 3 which consistently differs between host plant races of *T. conura*. This inversion is enriched for both metabolic processes and reproductive processes, providing a potential mechanism for coupling divergent ecological adaptation to reproductive isolation. We uncover that genes that are differentially expressed between feeding larvae of the different host races, the life stage under strongest viability selection, are enriched in the inversion. Finally, we document strong both pre- and postzygotic reproductive barriers, contributing to the potential for these host races to eventually form different species.

**Authors:** Runemark, Anna<sup>1\*</sup> Steward, Rachel<sup>1</sup> Ortega, Jesús<sup>2</sup> Nilsson, Kalle<sup>1</sup> Nolen, Zachary<sup>1</sup> Choudhary, Shruti<sup>3</sup> Yajuan Huang<sup>4</sup>

1. Department of Biology, Lund University, Sweden
2. Cavanilles Institute of Biodiversity and Evolutionary Biology, Spain
3. Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, Sweden
4. Department of Medical Biochemistry and Cell biology, Gothenburg University, Sweden

**Key words:** speciation, inversion, gene expression, host plant adaptation

**Email:** anna.runemark@biol.lu.se

**Abstract number:** O11

**Title:** Could selection for decreased variability in milk yield lead to asocial cows?

**Name:** Lars Rönnegård

**University:** SLU

**Abstract:**

Competition and indirect genetic effects (IGEs) result in a measurable genetic variation in variability, which can be estimated using a DHGLM producing estimated breeding values for variability (vEBV). We investigate how selection on vEBVs may affect the number of contacts a cow has in a model including IGEs. For each scenario, 20 replicates with 100 farms with 100 cows each were simulated. Milk yield was simulated using three correlated genetic components: direct, indirect and number of contacts. The number of contacts per cow was simulated using a social network algorithm. The simulated heritabilities were around 0.3 for direct and indirect genetic effects and 0.5 for number of contacts. The correlation between the vEBVs and the simulated true breeding values (TBVs) for the number of contacts was calculated. The results show that selection for reduced vEBVs can reduce the number of contacts a cow has.

**Authors:** Rönnegård, Lars Swedish University of Agricultural Sciences, Sweden  
Fikse, Freddy Växa, Swedish University of Agricultural Sciences, Sweden

**Key words:** phenotypic variability, social genetic effects, social network analysis, breeding value

**Email:** lars.ronnegard@slu.se

**Abstract number:** O12

**Title:** Evolution of flower shape in a coevolved system

**Name:** Pamela Santana

**University:** Lund University

**Abstract:**

Pollination interactions typically provide mutual fitness benefits to individuals of species interacting. Variation in the fitness benefits creates opportunities for natural selection to shape the traits mediating the interaction. Often, plant-pollinator interactions are mediated by multiple traits, and their combination ultimately determines the final interaction outcome. Plant-pollinator interactions face additional complexity, as change in one species may drive evolutionary change also in the interacting species, i.e. coevolution. The dynamic nature of coadaptation imposes a challenge for understanding how complex traits evolve in both species, and the reciprocity of evolution will depend also on the importance of the surrounding pollinator community. Here, we begin to address this topic from the plant perspective by investigating trait modularity and integration of *L. bolanderi* flowers, as flower modular organization is a key aspect to ensure evolvability. Our results indicate that coevolution is facilitated by trait modularity.

**Authors:** Pamela C. Santana<sup>1</sup>, Karin Gross<sup>1,2</sup>, John N Thompson<sup>3</sup>, and Magne Friberg<sup>1</sup>

<sup>1</sup> Department of Biology, Lund University, Sölvegatan 37, 223 62 Lund, Sweden <sup>2</sup> Department of Environment and Biodiversity, Paris Lodron University of Salzburg, Hellbrunnerstrasse 34, 5020 Salzburg, Austria <sup>3</sup> Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, Santa Cruz, CA 95060, USA

**Key words:** Pollination, Fitness, Trait integration, Modularity, Evolvability

**Email:** [pamela.santana@biol.lu.se](mailto:pamela.santana@biol.lu.se)

**Abstract number:** O13

**Title:** A novel supergene explains alternative reproductive strategies in the ant *M. ruginodis*

**Name:** Hanna Sigeman

**University:** Uppsala University

**Abstract:**

Many ant species have two alternative reproductive strategies. Large queens disperse to form single-queen colonies while small queens remain in their natal colonies as part of multi-queen groups. Although the genetic basis of these strategies has been elucidated in five ant lineages, many species remain unstudied. We investigate the genetic basis of alternative reproductive strategies in a novel system: the queen-size dimorphic ant *Myrmica ruginodis*. We captured 95 queens from 31 colonies in southern Finland, and using whole-genome sequencing, we found a novel 9 Mb supergene linked to queen size and social organization. Queens homozygous for the ancestral haplotype (“AA”) were larger and found only in single-queen colonies, while multi-queen colonies contained mixed haplotypes (“AB” and “BB”). This supergene is non-homologous to previously identified supergenes in ants, representing a novel evolutionary pathway to alternative reproductive strategies in ants.

**Authors:** Sigeman Hanna, University of Oulu (Finland) / Uppsala University (Sweden)

**Key words:** Supergene, genomics, ants, queen

**Email:** hanna.sigeman@oulu.fi

**Abstract number:** O14

**Title:** The role of introgression for adaptation of Atlantic herring to the Baltic sea

**Name:** Mafalda Sousa Ferreira

**University:** SciLifeLab Stockholm University

**Abstract:**

Introgressive hybridization can allow adaptation to new environments, but it can also impact fitness and compromise species survival. We study the role of genomic introgression for adaptation and divergence in two sister marine fish species, the Atlantic and Pacific herring, which are in contact in the Arctic Ocean. Using high-coverage whole genomes of 125 individuals representing the distribution of Atlantic and Pacific herring as well as the contact zone, we confirm previously described signatures of gene flow from Atlantic to Arctic Pacific herring, but find almost no introgression in the opposite direction. The remarkable exception is the spring-spawning Baltic herring ecotype, where we find Arctic Pacific introgression in high frequency in approximately 0.29% of the genome. Our analysis suggests that the introgressed genomic regions harbor genes involved in body growth and lipid metabolism and thus may have contributed to ecological adaptation in the Baltic Sea.

**Authors:** (1) Ferreira MS, Department of Zoology, Science for Life Laboratory, Stockholm University, Stockholm, Sweden (2) Felkel S, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden (3) Fuentes-Pardo AP, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden. Science for Life Laboratory, Data Centre. Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden (4) Goodall J, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden (5) Daniels RJ, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden (6) Pettersson M, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden (7) Andersson L, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden. Texas A&M University, Veterinary Integrative Biosciences, College Station, TX, USA

**Key words:** Adaptation, Introgression, Atlantic Herring, Genomics

**Email:** mafalda.ferreira@zoologi.su.se

**Abstract number:** O15

**Title:** Prenatal maternal stress underlies multi-hierarchical behavioural plasticity

**Name:** Diego Stingo Hirmas

**University:** Linköpings Universitet

**Abstract:**

Although prenatal stress (PS) has been observed to have long-term effects on the behaviour of many species, results vary significantly across studies. Recent research has shown that consistent differences in individual predictability exist when measured across a reaction norm. We aimed to assess whether PS could not only influence an animal's mean behavioural response but also its variability across multiple measurements. Hens of an advanced intercross of Red junglefowl and white leghorn chickens were implanted with either corticosterone to simulate maternal stress or placebo pellets subcutaneously. The offspring were repeatedly tested in an open-field test. We found that prenatal maternal corticosterone levels affected the offspring's mean anxiety-related response as well as how predictable they were in their behavioural responses. These results show that PS can cause heterogeneity of within-individual behavioural variability and highlight the role of reaction norms as a significant component when studying prenatal stressors' effect on animal behaviour.

**Authors:** Stingo-Hirmas Diego, Linköping University, Sweden. Rönnegård Lars, Swedish University of Agricultural Sciences Uppsala, Sweden. Wright Dominic, Linköping University, Sweden. Henriksen Rie, Linköping University, Sweden.

**Key words:** prenatal stress, behavioural plasticity, anxiety, predictability, residual intra-individual variation

**Email:** diego.stingo@liu.se



**Abstract number:** O16

**Title:** A study of the aging transcriptome from an adaptive perspective

**Name:** Zeeshan Syed

**University:** Linköping Unoversity

**Abstract:**

Differential gene-expression (DE) with age is generally attributed to the organism's reduced ability to maintain optimal expression. An alternative possibility is that a significant portion of DE is an adaptive response to poor physiological condition. Here, we test this hypothesis by genetically manipulating condition (independent of age) in female *Drosophila melanogaster*. Transcriptomic comparison of DE in the head due to reduced condition showed substantial overlap with that due to age. Gene ontology (GO) analyses revealed adaptive responses to poor condition and cellular damage by down- and up-regulated genes respectively. Further, downregulated (by either or both age and condition) genes belonging to the enriched GO terms showed (a) negatively correlation between tissue-specificity and the extent of DE and (b) signatures of purifying selection. These results suggest that a substantial proportion of the aging transcriptome is consistent with condition-dependent adaptive expression aimed at mitigating rather than causing aging.

**Authors:** Zeeshan A. Syed(1), Martin I. Brengdahl(1), Vinesh N. Shenoi(1), Avani Mital(1), Christopher M. Kimber(1), Helen Ekman(1), Malin Larsson(2), Claudio Mirabello(2), David Berger(3), Urban Friberg(1)

- (1) IFM Biology, Linköping University, 581 83 Linköping, Sweden
- (2) Department of Physics, Chemistry and Biology, National Bioinformatics Infrastructure Sweden, Science for Life Laboratory, Linköping University, 581 83 Linköping, Sweden
- (3) Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

**Key words:** aging, transcriptomics, condition-dependence, adaptation

**Email:** zeeshan.syed@liu.se

**Abstract number:** O17

**Title:** Testing genomic predictions of sexually antagonistic coevolution: Natural variation in the sex-peptide network in *Drosophila melanogaster*

**Name:** R. Axel W. Wiberg

**University:** Stockholm University

**Abstract:**

Sexually antagonistic coevolution (SAC) drives evolutionary dynamics, with males and females evolving traits that enhance their fitness but harming the opposite sex. While SAC is well-studied phenotypically, its genomic basis remains poorly understood. In *Drosophila melanogaster*, conflicts over re-mating rates is likely mediated by the sex-peptide (SP) network, a set of interacting male and female proteins. Males benefit from reducing female remating rates to ensure paternity, while females may should prefer to remate when advantageous. However, the role of natural genetic variation in the SP-network in shaping these behaviors is understudied. Here, we test genomic predictions of SAC in *D. melanogaster* populations identified by natural patterns of genetic co-variation at SP-network loci. We compare populations where males, females, or neither sex has an inferred advantage. We hypothesize that populations with a female advantage will show higher remating rates and reduced fitness costs to females compared to populations favoring males.

**Authors:** Thorson Alma, Sweden W. Wiberg Axel, Sweden R. Rhonda Snook, Sweden

**Key words:** Sexual conflict, sexually antagonistic coevolution, sex-peptide network

**Email:** axel.wiberg@zoologi.su.se

**Abstract number:** O18

**Title:** Dialectical relationships between microevolution and macroevolution

**Name:** Masahito Tsuboi

**University:** Lund University

**Abstract:**

Variation enables short-term evolution (microevolution), but its role in long-term evolution (macroevolution) remains debated. In my contribution, I will present analyses of extensive and multivariate *Drosophila* wing data spanning six levels of biological organization. Our results reveal consistent positive correlations between microevolutionary variation and macroevolutionary divergence, from variation within an individual to 40 million years of macroevolution. Surprisingly, the strongest relationship was observed between developmental noise and macroevolutionary divergence—levels thought to be the most distant—whereas the relationship between standing genetic variation and population divergence was modest, despite established theoretical predictions. Based on these results, I will propose pluralistic and dialectical relationships between microevolution and macroevolution, and discuss how data-driven approaches can integrate theory to advance 21st-century life sciences.

**Authors:** Tsuboi Masahito (Lund University, Sweden) Saito Keita (Chiba University, Japan) Takahashi Yuma (Chiba University, Japan)

**Key words:** evolutionary prediction, macroevolution, causation, data-and-theory-driven

**Email:** [masahito.tsuboi@biol.lu.se](mailto:masahito.tsuboi@biol.lu.se)

**Abstract number:** O19

**Title:** Domestication and Early Stress Resilience: Insights from White Leghorns and Red Junglefowl.

**Name:** Enya Van Poucke

**University:** Linköping University

**Abstract:**

Stress tolerance is a key factor in domestication, with studies showing that domestic species generally exhibit greater resilience than their wild ancestors. This study compares the stress responses of domestic White Leghorn (WL) laying hen chicks to those of their ancestral Red Junglefowl (RJF) when subjected to a pseudo-commercial hatchery procedure. Forty-four WL chicks and 40 RJF chicks underwent a series of stress-inducing treatments, including exposure to industrial noise and handling. Behavioral tests assessed their stress reactivity. Results indicated that RJF chicks emerged faster in a novel arena and vocalized more during social isolation, suggesting increased exploration and a stronger tendency to re-establish social contact. Interestingly, both WL and RJF chicks from the hatchery treatment showed lower corticosterone (CORT) levels during restraint tests compared to controls, hinting at potential stress-priming. Overall, domestication may not significantly affect stress resilience in this context, with adaptability possibly linked to their precocial nature.

**Authors:** Van Poucke Enya, IFM Biology, AVIAN Behavioural Physiology and Genomics Group, Linköping University, Sweden. Jensen Per, IFM Biology, AVIAN Behavioural Physiology and Genomics Group, Linköping University, Sweden.

**Key words:** Domestication, Early Stress, Commercial Hatchery

**Email:** enya.van.poucke@liu.se

**Abstract number:** O20

**Title:** Identifying the elusive migration genes

**Name:** Staffan Bensch

**University:** Lund University

**Abstract:**

It is amazing how tiny songbirds, weighing less than 10 grams, can fly with a strong determination in a specific direction, crossing huge geographical barriers to reach their winter grounds on another continent, all guided by their genes. Recently, we found a large and complex genomic region that directs willow warblers of two subspecies, trochilus and acredula, to different parts of Africa with great accuracy. We call this region MARB (Migration Associated Repeat Block). It seems to be passed down as a single unit without mixing, but we do not yet know if similar regions exist in other species or where exactly it is in the genome. In this talk, I will discuss the methods we have used to tackle these questions and share our progress in understanding how MARB influences songbird migration routes.

**Authors:** Staffan Bensch, Lund University, Sweden

**Key words:** Bird migration, genomics

**Email:** [staffan.bensch@biol.lu.se](mailto:staffan.bensch@biol.lu.se)

**Abstract number:** O21

**Title:** NA

**Name:** David Wheatcroft

**University:** Stockholms universitet

**Abstract:**

Population divergence in mating traits and preferences plays a critical role in speciation. Thus, uncovering how population divergence emerges and how it is maintained is a long-term goal in evolutionary biology. Songbird song is acquired through social learning, which has alternatively been hypothesized to facilitate or impede population and species differences. Here, I outline my group's recent work on song divergence across pied flycatcher (*Ficedula hypoleuca*) populations and exploring the implications for reproductive isolation with the closely related collared flycatcher (*F. albicollis*) where they co-occur. Our approach combines field experiments, acoustic analyses, RNA sequencing, and captive rearing.

**Authors:** Wheatcroft David, Stockholm University, Sweden

**Key words:** Songbirds, mating traits, speciation

**Email:** david.wheatcroft@zoologi.su.se

**Abstract number:** O22

**Title:** Strong Selection, but low repeatability: Temperature-specific effects on genomic predictions of adaptation

**Name:** David Berger

**University:** Uppsala University

**Abstract:**

Climate warming is increasing temperatures beyond the optima of many ectotherms, which imposes strong selection on basal physiological and metabolic processes. This suggests that adaptation to climate warming should be relatively predictable. We tested this hypothesis from the level of genes to life-history traits by conducting an evolve-and-resequence experiment in a cosmopolitan insect pest. Indeed, phenotypic evolution was faster and more repeatable at hot temperature. However, at the genome-level, adaptation to heat was less repeatable than cold-adaptation. Moreover, while genomic predictions of phenotypic (mal)adaptation to hot temperature were accurate when applied to populations of the same geographic origin, they were highly inaccurate for populations of different geographic origin. These results seem best explained by an increased importance of epistasis during adaptation to heat and imply that genomic predictions of adaptation in key life-history traits, which mediate the agricultural impact of insect pests, may become increasingly difficult as climate warms.

**Authors:** Rego, Alexandre Uppsala University, Sweden

Julian Baur Uppsala University, Sweden

Camille Girard-Tercieux Uppsala University, Sweden

Maria de la Paz Celorio-Mancera Stockholm University, Sweden

Rike Stelkens Stockholm University, Sweden

David Berger Uppsala University, Sweden

**Key words:** Repeatability of evolution, climate change, pest impact

**Email:** david.berger@ebc.uu.se

**Abstract number:** O23

**Title:** Contrasting evolution in incipient versus established species

**Name:** Folmer Bokma

**University:** Karlstad University

**Abstract:**

Are differences between species the long-term consequence of microevolution within species, or does speciation involve fundamentally different processes? I developed a phylogenetic method that decomposes interspecific trait covariance into two components: trait coevolution along phylogeny branches and at bifurcations. Applying this method shows, for example, that primate brain and body size evolve less correlatedly during speciation than between speciation events. With improved computational methods, we could apply this approach to large trait matrices, like gene expression data, to explore the molecular basis of the tempo and mode in phenotypic evolution. This opens new avenues for data-driven study of speciation, evolvability, and the relation between micro- and macroevolution.

**Authors:** Folmer Bokma

**Key words:** speciation, macroevolution, trait covariance, gene expression, coevolution

**Email:** folmer.bokma@kau.se



**Abstract number:** O24

**Title:** How ecologically relevant resource distributions influence condition-dependent trade-offs

**Name:** Lennart Winkler

**University:** SLU Uppsala

**Abstract:**

Trade-offs are a central part of evolutionary biology. Individuals are assumed to trade investment in one energetically demanding trait against another, particularly when resources are limited. Most previous studies experimentally tested for trade-offs by manipulating resources quantity (e.g., high vs. restricted resources). However, this approach is unlikely to reflect the distribution of resources in natural populations. Here, we examine how a range of ecologically relevant distributions of resources influences trade-offs among pre-copulatory, post-copulatory and immune traits in male field crickets. Our results indicate that the emergence of trade-offs was sensitive to the underlying distribution of resources among males. Importantly, trade-offs were more likely when only high and low condition individuals were included – the least ecologically relevant resource distribution we considered. This cautions that experimental designs that do not reflect ecologically relevant conditions might identify trade-offs that are less relevant in natural populations.

**Authors:** Winkler, Lennart, Stockholm University, Sweden Simmons, Leigh, University of Western Australia, Australia Fitzpatrick, John, Stockholm University, Sweden

**Key words:** Trade-offs, Condition-dependence, Sexual Selection, Evolutionary Ecology

**Email:** [lennart.winkler@zoologi.su.se](mailto:lennart.winkler@zoologi.su.se)

**Abstract number:** O25

**Title:** From Wild to Tamed and Back: the evolutionary history of Sardinian sheep

**Name:** Marianne Dehasque

**University:** Uppsala University

**Abstract:**

Sardinia hosts three sympatric sheep lineages with distinct selective histories: the Sardinian mouflon, likely representing feral remnants of early domesticated sheep

**Authors:** Dehasque Marianne. Uppsala University, Sweden

**Key words:** the Nera sheep, a local breed with ancestral traits

**Email:** marianne.dehasque@ebc.uu.se

**Abstract number:** O26

**Title:** Why All PCAs in Evolutionary Biology are Likely Wrong

**Name:** Eran Elhaik

**University:** Lund University

**Abstract:**

Irreproducibility in science hinders progress, wastes resources, and poses risks and biases. Addressing this issue requires examining commonly used methods, such as principal components analysis (PCA), a popular statistical tool in evolutionary biology and related fields. Despite its widespread use and citation, PCA's accuracy for genetic applications has never been conclusively proven. We argue that its results are often artifacts, and its popularity stems from the ease of manipulating outcomes, creating significant reproducibility issues. This is particularly problematic in ancient DNA studies, where bioinformatics methods are critical. To support our claims, we developed a model that compares true genetic distances to those produced by PCA, showing its bias and propensity to generate misleading results. Using real genetic data, we demonstrate how PCA leads to false conclusions and propose an alternative approach for measuring genetic distances. Our findings suggest PCA is unsuitable for genetic studies and call for reevaluation ~250,000 studies.

**Authors:** Eran Elhaik, Lund University, Sweden

**Key words:** PCA, evolutionary biology, paleogenomics, physical anthropology, reproducibility crisis

**Email:** eranelhaik@gmail.com

**Abstract number:** O27

**Title:** Investigating the role of meiotic recombination in social insects - Termites adding an important piece to the puzzle

**Name:** Turid Everitt

**University:** Uppsala University

**Abstract:**

Meiotic recombination is an essential process in sexually reproducing organisms as it increases the efficiency of selection. There is however great variation in recombination rate across the tree of life, with eusocial insects from the order Hymenoptera in the topmost range. It has been hypothesized that eusociality leads to selection for high recombination rate as this would increase the genetic diversity in colonies and reduce kin conflict. The association between eusociality and high recombination rate has however not previously been studied outside of Hymenoptera. In this study we show that two termite species, which are eusocial insects from a different order, have substantially lower recombination rates than eusocial hymenopteran insects, contradicting aforementioned hypothesis. We present an alternative hypothesis, relating the high recombination rate in eusocial hymenopteran insects to their skewed sex-ratio of reproductive individuals, where the high number of haploid males present an efficient way to test different haplotypes.

**Authors:** Everitt Turid, Uppsala University, Sweden Rönneburg Tilman, Uppsala University, Sweden Elsner Daniel, University of Freiburg, Germany Olsson Anna, Uppsala University, Sweden Liu Yuanzhen, Uppsala University, Sweden Larva Tuuli, Uppsala University, Sweden Korb Judith, University of Freiburg, Germany Webster Matthew, Uppsala University, Sweden

**Key words:** Recombination rate, eusociality, selection, genomics

**Email:** turid.everitt@imbim.uu.se

**Abstract number:** O28

**Title:** The evolution of conspicuous markings in skates and rays

**Name:** John Fitzpatrick

**University:** Stockholm University

**Abstract:**

Prey use a wide range of anti-predator defences to avoid predation. Yet few studies have examined the evolution of multiple anti-predator defences while simultaneously incorporating ecologically relevant factors at a macroevolutionary scale. Here we examine the evolution of anti-predator defences in 580 skates and rays (>90% of extant species), a diverse group of cartilaginous fishes. We demonstrate that skates and rays have evolved distinct anti-predator defences that are shaped by ecological factors. In particular, skates and rays lacking robust anti-predator defences and living at shallower depths were more likely to evolve conspicuous anti-predator markings than species with robust defences and species living at deeper depths. Together our findings suggest that the evolution of conspicuous markings is predominantly driven by trade-offs among alternative anti-predator defences and the visual environment in skates and rays. These findings have wider implications for our understanding of functional trade-offs in anti-predator defences.

**Authors:** Madicken Åkerman, Ana C.R. Gomes, David Wheatcroft, Niclas Kolm, Karl Gotthard and John L. Fitzpatrick

Department of Zoology, Stockholm University, Svante Arrhenius väg 18b, 106 91 Stockholm, Sweden

**Key words:** predation, visual signals, trade-offs, defence portfolio

**Email:** john.fitzpatrick@zoologi.su.se

**Abstract number:** O29

**Title:** Step by step, an adaptive radiation is made: Empirical evidence for the 'least action effect'

**Name:** José Cerca

**University:** Naturhistoriska

**Abstract:**

Abs: Adaptive radiations occur when a single lineage diversifies to exploit a broad range of ecological niches. However, the sequence of ecological occupation remains poorly understood. Two potential scenarios are herein proposed: random occupation, where species fill niches without a discernible pattern, and progressive occupation, where species track specific environmental gradients in a systematic manner. In this study, I examined four well-known adaptive radiations: Galápagos finches (Galápagos), *Descurainia* (Macaronesia), *Aeonium* (Macaronesia), and Anoles (West Indies). By analyzing phylogenetic and ecological distances between species, I found evidence supporting the hypothesis of progressive ecological occupation. This suggests that species diversify along a structured niche axis, implying that adaptive radiations may follow a more predictable, directional pattern. These findings are framed within the context of the 'least action principle' described by Gavrilets and Vose (2005).

**Authors:** None

**Key words:** Adaptive radiation

**Email:** jose.cerca@gmail.com

**Abstract number:** O30

**Title:** Experimental examination of the neural background to cognitive variation across species in Poeciliid fishes

**Name:** Ana Gomes

**University:** Stockholm University

**Abstract:**

Cognitive performance is hypothesized to be linked to brain size. However, the few phylogenetically controlled studies that have tested this hypothesis focus primarily on mammals and birds, consider only distantly related species, and have not examined individual brain regions. Here, we overcome these limitations by experimentally quantifying cognitive performance and brain size in 173 individuals from 18 fish species from a single family of fishes - the Poeciliidae. For all individuals, we assessed performance on three cognitive tasks: reversal learning, detour-reaching, and object permanence. We then measured each individual's total brain volume, and the volume of different brain regions. This approach offers a robust assessment of the relation between relative brain volume and cognition. Indeed we find evidence that brain regions may be important in some cognition axis. Thus our results shed light on the evolution of cognitive performance and highlight the importance of considering the scale of macroevolutionary studies.

**Authors:** Gomes, Ana Cristina R., De Meester, Gilles, Boussard, Annika, Shamsgovara, Arezo, Amcoff, Mirjam, Wheatcroft, David, Kolm, Niclas, Fitzpatrick, John L.

Department of Zoology/Ethology, Stockholm University, Svante Arrhenius Väg 18B, SE-10691 Stockholm, Sweden

**Key words:** cognition, brain size, brain regions, Poeciliidae, fish

**Email:** ana.gomes@zoologi.su.se

**Abstract number:** O31

**Title:** Atlantic Herring - Population trends and perspectives from the Baltic Sea

**Name:** Jake Goodall

**University:** Uppsala University

**Abstract:**

The Atlantic herring (*Clupea harengus*) is one of the most abundant vertebrates on earth, with an effective population size of one trillion. The species exhibits minute genetic differentiation at neutral loci and strong differentiation across selectively adaptive loci underlying ecological adaptation. Recently, a novel herring-specific SNP array was developed, increasing the accessibility of large-scale population genetic studies for the species. Here, we present ongoing research from the Baltic Sea, a region experiencing significant fishing pressure across the majority of stocks. Notably, we aim to discuss ecosystem-scale clines in genetic variation across the Baltic relative to factors such as latitude and salinity. Additionally, we discuss the discovery of novel Baltic ecotypes, such as the recently described Slåttersill, a large, piscivorous Baltic herring. Overall, we aim to highlight novel herring research and explore variable patterns of genetic diversity found in the Baltic Sea.

**Authors:** Goodall, J., Uppsala University, Sweden. Pettersson, M., Uppsala University, Sweden. Cocco, A., Uppsala University, Sweden. Laikre, L., Stockholm University, Sweden. Andersson, L., Uppsala University, Sweden

**Key words:** Herring, Population Genetics, SNP, Ecotype

**Email:** jake.goodall@imbim.uu.se



**Abstract number:** O32

**Title:** Adaptive benefits of hybridization in *Saccharomyces* yeast under thermal stress: genomic and fitness analyses

**Name:** Chloé Haberkorn

**University:** Stockholm University

**Abstract:**

Accelerating climate change urges us to better understand and predict evolutionary responses to temperature shifts. Hybridization, by increasing genetic variation, can widen the range of adaptive responses and genetic mechanisms available to populations, which may allow them to survive fluctuating temperature conditions. Here, we explored the evolutionary consequences of hybridization between two species of *Saccharomyces* yeast, by evolving both hybrid and parental strains for 200 generations under hot, cold and fluctuating temperature regimes, and collecting whole genome sequencing data. Fitness was measured for each population, in all three temperature conditions, before and after experimental evolution. Overall, hybrids showed intermediate fitness to their parents, but a few hybrids evolved in hot conditions were clearly winning, with a higher fitness than that of their ancestors, in all three temperatures. Genomic exploration of mutations in hot-evolved hybrid populations revealed that hybridization can affect cellular stress-management pathways, leading to increased fitness in hot environments.

**Authors:** Chloé Haberkorn, Gerda Saxer, Noah Gettle, Jacob Elsen, Nahui Medina Chavez, Duncan Greig, Michael Travisano, Rike Stelkens

**Key words:** experimental evolution, climate change, yeasts, hybridization

**Email:** [chloe.haberkorn@zoologi.su.se](mailto:chloe.haberkorn@zoologi.su.se)

**Abstract number:** O33

**Title:** The importance of standing ancestral variation in behavioural evolution

**Name:** Christina Hansen Wheat

**University:** Stockholm University

**Abstract:**

Understanding how behaviours evolve is a long-standing goal in evolutionary biology. While novel mutations can be strong drivers of phenotypic change, a growing body of evidence now suggests that evolutionary change relies heavily upon standing genetic variation. However, the highly complex genetic basis of behavioural traits makes it challenging to establish the ancestral origins of behaviour through genomic studies. An alternative approach is to compare behaviours between groups of ancestral and derived species. In this context, I will use examples from separate behavioural tests conducted on European grey wolves and domestic dogs hand-raised under identical conditions. The results highlight two key points: 1) two behaviours (i.e. fetching and human-directed attachment) widely labelled as being unique to dogs are expressed in wolves, suggesting that selection acts on standing ancestral variation for these traits, and 2) identifying rare behavioural variation among ancestral populations could provide significant insights into how behaviours evolve.

**Authors:** Hansen Wheat, Christina

Stockholm University, Sweden

**Key words:** Behavioural evolution, standing variation

**Email:** christina.hansen@zoologi.su.se

**Abstract number:** O34

**Title:** Epigenetic regulation of immune response in the Swedish sand lizard (*Lacerta agilis*)

**Name:** Joshua Hufton

**University:** Uppsala universitet

**Abstract:**

Sand lizards (*Lacerta agilis*) are found across Europe, even stretching as far north to extremely cold environments such as in Sweden. Being highly adapted to such extreme temperatures makes them highly susceptible to climate change, and the knock-on effects this could have on disease dynamics. Using an experimentally isolated, outbred, island population in southern Sweden we are investigating immunogenetics and epigenetics in these highly adaptable populations. Using immune challenge experiments, we are characterising the sand lizard immune response using RNA and bisulfite sequencing, and assessing the role of DNA methylation in reptilian immunity. This study will greatly improve our understanding of epigenetic regulation of immunity in reptiles in general but also give us key insights into specific genes underlying the Swedish sand lizard immune response. Our results are also important for the conservation of the species in Sweden, which are expected to face novel changes in disease dynamics as a result of climate change.

**Authors:** Hufton Joshua, Institutionen för ekologi och genetik, Uppsala Universitet, Sweden, Olsson Mats, Institutionen för biologi och miljövetenskap, Göteborgs universitet, Sweden Wapstra Erik, School of Natural Sciences, University of Tasmania, Tasmania, Lillie Mette, Institutionen för ekologi och genetik, Uppsala Universitet, Sweden

**Key words:** Epigenetics, reptiles, immunity, gene expression, conservation

**Email:** [joshua.hufton@ebc.uu.se](mailto:joshua.hufton@ebc.uu.se)

**Abstract number:** O35

**Title:** The SciLifeLab Planetary Biology Capability

**Name:** Jacob Höglund

**University:** Uppsala

**Abstract:**

Planetary Biology Capability SciLifeLab's Planetary Biology capability sprung out of a need for trans-disciplinary and coordinated approaches to study life on Earth, with a broad scope ranging from single molecules and cells to individual species, species communities, ecosystems, and their functioning on the planet. Simply speaking "Life in environmental context". By combining SciLifeLab infrastructures, and the data-driven life science approach, the capability will create opportunities with a strong impact on biodiversity oriented research (broadly defined). Taking advantage of SciLifeLab's national research infrastructure, technology, and scientific excellence, techniques such as imaging, microscopy, genomics, bioinformatics, metabolomics, proteomics, and big data analysis, for non-biomedical research will be used to shed light on all aspects of life on our planet.

**Authors:** Höglund Jacob, Uppsala University, Sweden

**Key words:** SciLife, Planetary Biology, Biodiversity, Capability, Platforms

**Email:** jacob.hoglund@ebc.uu.se

**Abstract number:** O36

**Title:** Characterisation and quantification of deleterious genetic variants in non-model organisms: from present to extinct species

**Name:** Julia Höglund

**University:** Stockholm University

**Abstract:**

Many animal populations are currently undergoing dramatic decline

**Authors:** Julia Höglund<sup>1,2</sup>, Seyan Hu<sup>1</sup>, Martijn Derks<sup>1</sup>, Love Dalén<sup>2</sup>, Mirte Bosse<sup>1,3</sup>

<sup>1</sup> Dept. of Animal Breeding and Genomics, Wageningen University & Research, Wageningen, THE NETHERLANDS <sup>2</sup> Centre for Palaeogenetics, Dept. of Zoology, Stockholm University, Stockholm, SWEDEN <sup>3</sup> Section Ecology & Evolution, Amsterdam Institute for Life and Environment (A-LIFE), Vrije Universiteit Amsterdam, THE NETHERLANDS

**Key words:** when a population becomes smaller, threats affecting the species' genome become larger. A reduced population size increases the risk of inbreeding and loss of beneficial genetic variation, results in a worse ability to adapt to changes, and can potentially lead to extinction. Hence, there is a critical need to quantify damaging genetic variation to aid conservation efforts and understand past extinctions. By expanding previous species-specific models scoring genetic variants based on predicted deleteriousness, we created a model that is able to predict deleteriousness also across non-model organism. We trained the initial model using deep sequencing data from domesticated pig, and our next step is to compare it to a model trained on wild boar, using the same pipeline. By extending beyond model species, we enhance its applicability to several species, hopefully providing insights into the possible genetic factors underlying species decline.

**Email:** julia.hoglund@su.se

**Abstract number:** O37

**Title:** Loss of genetic variation and establishment of damaging variants in effectively small populations

**Name:** Martin Johnsson

**University:** Swedish University of Agricultural Sciences

**Abstract:**

Domestic animal populations, as well as many animal populations in the wild, are effectively small and shrinking. In particular, in domestic animals with breeding programs, there is very strong directional selection, while breeds without modern breeding tend to be fragmented and in decline. Large-effect deleterious variants at relatively high frequencies are common enough in domestic animals that they pose applied problems for breeders. In our work, we have combined genomic data analysis with a series of population genetic models to address the loss of genetic variation in effectively small and shrinking populations, and how large-effect deleterious variants establish themselves in small populations. The results suggest the presence of transient genetic variation, predicted to be gradually lost, and that balancing selection by heterozygote advantage, even temporarily through close linkage, makes establishment of new large-effect deleterious alleles likely.

**Authors:** Martin Johnsson

Department of Animal Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden.

**Key words:** domestic animals, population genetics, deleterious variation

**Email:** martin.johnsson@slu.se

**Abstract number:** O38

**Title:** The adaptive evolution of herring to different salinities

**Name:** Cheng Ma

**University:** Uppsala Universitet

**Abstract:**

Understanding the relationship between genetic and phenotypic variation is a fundamental goal in evolutionary biology. For marine organisms, salinity is a crucial environmental factor affecting survival and adaptation. However, the genetic mechanisms driving adaptive evolution to different salinity levels remain largely unexplored. Herring, which are widely distributed across high-salinity Atlantic Ocean and low-salinity Baltic Sea environments, serve as an ideal model for investigating salinity adaptation. Our research reveals that copy number variation (CNV) in hatching enzyme genes supports herring adaptation to diverse salinity conditions. Specifically, we discovered a significant expansion in the copy number of a gene encoding a hatching enzyme on chromosome 26 in Baltic herring. We hypothesize that this expansion is a selective response to the low salinity of the Baltic Sea, which likely reduces enzyme activity relative to the Atlantic's marine conditions. This CNV exemplifies an evolutionary mechanism that enhances herring survival and reproductive success.

**Authors:** Cheng Ma, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden.

Fahime Mohamadnejad Sangdehi, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden.

Mats Pettersson, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden.

Leif Andersson, Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden.

**Key words:** Adaptive evolution, salinity, herring, hatching enzyme, CNV

**Email:** cheng.ma@imbim.uu.se

**Abstract number:** O39

**Title:** Toward predictive evo-devo: a data-driven approach to modeling phenotypic variation

**Name:** Lisandro Milocco

**University:** Stockholm University, SciLifeLab

**Abstract:**

Predicting evolution is crucial for tackling challenges like population survival under climate change and controlling emerging pathogens. However, it remains limited due to the difficulty in predicting the appearance of new phenotypic variation, generated by the notoriously complex process of development. Existing approaches to model phenotypic variation fall into two main categories: statistical models, which treat development as a black box—describing but not predicting variation—and mechanistic models, which require detailed developmental information that is rarely available. In this talk, I will present a data-driven framework that addresses these limitations by modeling the emergence of variation through time delay embedding of the developing system's states. I will share preliminary *in silico* results using a gene regulatory network and discuss ongoing efforts to apply this framework to empirical datasets, highlighting its potential to enhance our ability to predict evolution.

**Authors:** Milocco, Lisandro. Stockholm University, SciLifeLab. Sweden.

**Key words:** evo-devo, prediction, development, time series

**Email:** milocco.lisandro@gmail.com



**Abstract number:** O40

**Title:** Speciation and Temporal Dynamics of Female Sperm Management

**Name:** Dick Moberg

**University:** Stockholm University

**Abstract:**

Understanding how traits cause reproductive isolation between diversifying lineages is a central focus of evolutionary biology. However, mechanisms causing reproductive isolation early in the speciation process are seldom studied since research has primarily focused on already formed species in the late stages of speciation. In this study, we investigate mechanisms causing early post-mating prezygotic (PMPZ) reproductive isolation between two populations of *Drosophila montana*. In this species, crosses between populations results in reduced fertilization. We therefore hypothesized that heteropopulation crosses would exhibit different rates of sperm utilization compared to conpopulation crosses. Our findings provide partial support for this hypothesis. We observed significant differences in sperm storage dynamics in one heteropopulation cross, which exhibited a slower rate of filling their sperm storage organs and retaining sperm for a longer period compared to other crosses. This suggests that inefficient sperm utilization may be a mechanism of PMPZ reproductive isolation, potentially influenced by mismatches in sperm and reproductive tract lengths or differences in seminal fluid composition.

**Authors:** Moberg Dick, Stockholm University, Sweden Erin Macartney, University of Sydney, Australia Mathilda, Pembury-Smith, Stockholm University, Sweden Alma Thorsson, Stockholm University, Sweden Rhonda Snook, Stockholm University Sweden

**Key words:** Speciation, Reproductive Isolation, postcopulatory mechanisms

**Email:** dick.moberg@zoologi.su.se

**Abstract number:** P1

**Title:** The evolution of thermal performance curves across the model genus yeast

**Name:** Jennifer Molinet

**University:** Stockholm University

**Abstract:**

Accelerating climate change and extreme temperatures highlight the urgent need to understand the evolutionary impacts on biodiversity to develop conservation strategies for at-risk species. This study assesses heat stress adaptation across eight *Saccharomyces* species with varying temperature preferences sourced from diverse global habitats. A long-term evolution experiment with gradual temperature increases (25-40°C) was conducted to simulate climate change. The results showed a ~30% fitness improvement at higher temperatures in evolved populations. While cold-tolerant species had greater fitness gains, they did not reach warm-tolerant species' limits. Evolutionary trajectories of TPCs varied, with some species showing a trade-off between thermal breadth and upper thermal limit, while others increased their optimal temperature and thermal tolerance. Two main trajectories emerged: broader TPCs for genotypes with higher thermal optima and increased CTmax at the cost of maximum fitness. These findings emphasize the complexity of microbial climate responses and the need to consider species diversity.

**Authors:** Molinet Jennifer, Stockholm University, Sweden Stelkens Rike, Stockholm University, Sweden

**Key words:** *Saccharomyces*, thermal performance curve, thermal adaptation, experimental evolution, climate change

**Email:** jennifer.molinet@zoologi.su.se

**Abstract number:** P2

**Title:** The tempo and mode of toxicant sensitivity evolution

**Name:** Iain Moodie

**University:** Lund University

**Abstract:**

Chemical pollution is a critical driver of biodiversity loss, impacting ecosystems worldwide. Species differ remarkably in their sensitivities to environmental toxicants, and understanding why is crucial to better predict ecological risks and inform conservation strategies. By considering effect concentrations (the concentration of toxicant X that produces effect Y in species Z) as quantitative traits in a macro-evolutionary perspective, we can investigate how toxicant sensitivity evolves across diverse taxa. Here, we use large ecotoxicological databases of standardised effect concentrations (>1 million empirical results) for a wide variety of toxicants and species. Combined with time-calibrated molecular phylogenies, we use a data-driven phylogenetic comparative framework to elucidate the processes and patterns that have given rise to contemporary distributions of species' sensitivities. Our analysis revealed that the tempo and mode of toxicant sensitivity evolution differed across classes of toxicants, and recovered cases of convergent evolution across deeply diverged taxa.

**Authors:** Moodie, Iain R., Department of Biology, Lund University, Lund, Sweden De Lisle, Stephen P., Department of Environmental and Life Science, Karlstad University, Karlstad, Sweden

**Key words:** macroevolution, ecotoxicology, PCM (phylogenetic comparative methods)

**Email:** iain.moodie@biol.lu.se

**Abstract number:** P3

**Title:** Chromosomal rearrangements in Ithomiini butterflies

**Name:** Karin Näsval

**University:** Wellcome Sanger Institute

**Abstract:**

Chromosomal rearrangements can act as barriers to gene flow, promoting reproductive isolation and divergence between closely related species. However, their impact on speciation rates and importance in a macroevolutionary perspective remains unclear. Here, we focus on the South American Ithomiini butterflies, a diverse group of over 400 species with varying speciation rates among genera and highly variable karyotypes ( $n=5-120$ ). We aim to develop high-quality chromosome-level assemblies for 200 taxa across the Ithomiini phylogeny to investigate the relationship between chromosomal rearrangements and speciation rates. Preliminary findings reveal that rearrangements are more extensive than previous chromosome counts indicated, primarily involving inter-chromosomal fusions and fissions. We also observed significant variation in genome size correlated with transposable element expansion. In addition, we explore how chromosomal rearrangements arise and increase in frequency by examining the relationship between rearrangement rates, genetic diversity, and transposable element dynamics.

**Authors:** Näsval Karin, Tree of Life, Wellcome Sanger Institute, UK Salazar-Carrión Patricio, Department of Zoology, University of Cambridge, UK Bacquet Caroline, Universidad Regional Amazónica Ikiam, Tena, Ecuador Elias Marianne, Institut de Systématique, Evolution, Biodiversité, MNHN-CNRS-Sorbonne Université-EPHE-PSL-Université des Antilles, Muséum national d'Histoire naturelle, France. Willmott Keith, Florida Museum of Natural History, Florida, USA Salazar Camilo, Faculty of Natural Sciences, Universidad del Rosario, Bogota, Colombia. Meier Joana, Tree of Life, Wellcome Sanger Institute, UK

**Key words:** Speciation genomics, chromosomal evolution, structural variation, genome architecture

**Email:** karin.nasvall@sanger.ac.uk

**Abstract number:** P4

**Title:** The seasonal clock of animals: gaining insights via a molecular evolutionary analysis

**Name:** Saurav Baral

**University:** Stockholm University

**Abstract:**

The daily and seasonal cycles of all animals are controlled by their circadian clock, a dynamic network of interacting genes that keeps time. These clock genes, which are regulated by a complex transcription-translation feedback loop, are essential for synchronizing animal life cycles with favorable environmental conditions, such as resource availability. Unfortunately, for many animals, their circadian clocks are becoming desynchronized with resource availability due to anthropogenic factors such as light pollution and climate change. Worsening the issue is the limited understanding of how the circadian clock functions and evolves in most species. Here, I have analyzed clock genes from over a hundred different butterfly species to show that these genes are evolving at a far higher rate than expected. Although this is a conserved system present across all animals, it is also rife with adaptive evolution, altering complex interactions between different clock genes.

**Authors:** Saurav Baral, Stockholm University Christopher Wheat, Stockholm University

**Key words:** Clock gene, Molecular Evolution, Seasonal adaptaion

**Email:** baral.saurav@zoologi.su.se

**Abstract number:** P5

**Title:** Play ontogeny in young chickens - effects of domestication, tameness, sex, and the cognitive impact of play stimulation

**Name:** Rebecca Oscarsson

**University:** Linköping University

**Abstract:**

It is known that young chickens display different play behaviours, but the function of these are not fully understood. Moreover, domestication is believed to have increased playfulness, and tameness is hypothesized to be the main trait driving the domesticated phenotype. Furthermore, potential sex differences in play had not been investigated in chickens. Therefore, in three separate studies, we aimed to describe potential effects of (1) domestication, (2) tameness, and (3) sex, on chicken play ontogeny. First, we compared the ancestral Red Junglefowl (RJF) and the layer hybrid of White Leghorn (WL). Secondly, we compared Red Junglefowls selected for high (RJF HF) and low (RJF LF) fear of humans. Lastly, male, and female WL were compared. Finally, in a fourth study, the effects of play stimulation on the cognitive capacities of RJF and WL were explored. The results of these projects will be presented, and the function of play discussed.

**Authors:** Rebecca Oscarsson\* & Per Jensen IFM Biology, AVIAN Behaviour Physiology and Genomics Group, Linköping University, Linköping, Sweden  
\*Presenting author

**Key words:** Play behaviour, domestication, tameness, sex differences, function of play

**Email:** rebecca.oskarsson@liu.se

**Abstract number:** P6

**Title:** Genomic Erosion Through the Lens of Comparative Genomics

**Name:** Carolina Pacheco

**University:** Lund University

**Abstract:**

Comparative genomic analyses provide key insights for conservation by revealing genomic erosion patterns across species with different evolutionary histories. Understanding the mechanisms driving genomic erosion is crucial for species like the Mauritius parakeet, Mauritius kestrel, and pink pigeon, where conservation efforts led to demographic recovery, but genetic diversity keeps declining. Comparing the genomes of 39 bird species we found that species with larger historical effective population sizes have greater heterozygosity, but carry higher masked load, highlighting the importance of historical demography to assess species vulnerability to genomic erosion. We identified significant differences in realized load between taxonomic groups of our target species, possibly due to differences in life history traits. Structural variation analyses revealed differences in transposable elements and genomic rearrangements between groups. Our findings underscore the value of multispecies comparisons in understanding the evolutionary dynamics of genomic erosion and its relevance for biodiversity conservation.

**Authors:** Xuejing Wang, *Globe Institute, University of Copenhagen, Denmark*  
Claudia Fontseré, *Globe Institute, University of Copenhagen, Denmark*  
Cock Van Oosterhout, *School of Environmental Sciences, University of East Anglia, UK*  
Jim Groombridge, *Durrell Institute of Conservation and Ecology, University of Kent, UK*

Bengt Hansson, *Department of Biology, Lund University, Sweden*

Ximena Alva Caballero, *Department of Biology, Lund University, Sweden*  
Carolina Pacheco, *Department of Biology, Lund University, Sweden*  
Hernán E. Morales, *Globe Institute, University of Copenhagen, Denmark*  
\* These authors contributed equally.

**Key words:** Genomic erosion, comparative genomics

**Email:** carolina.pacheco@biol.lu.se

**Abstract number:** P7

**Title:** Have sponges stolen gene to survive without oxygen?

**Name:** Sofia Paraskevopoulou

**University:** Lund University

**Abstract:**

In animals, new phenotypic traits often emerge through gene duplications, splice variation, recombination, or even gene loss. Lateral Gene Transfer (LGT), the movement of genetic material across species borders, is another potential mechanism but is rare in animals due to constraints on vertical transfer of foreign DNA to offspring. Preliminary data from our lab suggests a potential LGT event in freshwater sponges, where the RQUA gene, likely transferred from anaerobic microbes, enables the synthesis of an anaerobiosis-specific cofactor rhodoquinone (RQ). This is unexpected, as no other animal encodes RQUA and those animals that can synthesize RQ use alternative splicing to yield a new gene product of an unrelated protein COQ2. Phylogenetic analysis suggests that freshwater sponges acquired RQUA via LGT, possibly from an euglenid protist. While RQUA expression in microbes is regulated by environmental oxygen levels, we observed subtle gene expression changes in sponges under varying oxygen conditions. Given that sponges experience tissue anoxia during contractions and asexual reproduction, we hypothesize that tissue hypoxia may drive RQ upregulation.

**Authors:** 1. Paraskevopoulou Sofia, Molecular Biosciences, Department of Biology, Lund University, Sweden 2. Vizitiu Mara, Molecular Biosciences, Department of Biology, Lund University, Sweden 3. Aguilera Campos Karla Iveth, Molecular Biosciences, Department of Biology, Lund University, Sweden 4. Itriago Humberto, Molecular Biosciences, Department of Biology, Lund University, Sweden 5. Boisard Julie, Molecular Biosciences, Department of Biology, Lund University, Sweden 6. Pereira Raquel, Molecular Biosciences, Department of Biology, Lund University, Sweden 7. Carr Page, Gonzaga University, USA 8. Boercker Lily, Gonzaga University, USA 9. Flandrin Laura, CNRS-Grenoble University, France 10. Pierrel Fabien, CNRS-Grenoble University, France 11. Shepherd Jennifer, Gonzaga University, USA 12. Horton April, Bates College, USA 13. Leys Sally, University of Alberta, Canada 14. Stairs Courtney, Molecular Biosciences, Department of Biology, Lund University, Sweden

**Key words:** gene expression, transcriptomics, confocal microscopy, sponge phenotype

**Email:** sofia.paraskevopoulou@biol.lu.se



**Abstract number:** P8

**Title:** Elevated temperature changes behaviour, selection and mating system landscapes

**Name:** Matilda Pembury Smith

**University:** Stockholm University

**Abstract:**

Climate change-induced heat stress can impair fertility and reproductive behaviour in both sexes, with intensified effects in monogamous species due to the risk of an exclusive pairing with a non-fertile individual. Shifts in mate choice and selection for polyandry have been theorised as population fitness rescue mechanisms under climate change. However, the impact of heat stress on mating behaviour in both sexes remains poorly understood. Here, we tested how elevated developmental temperature influences reproductive behaviour using isofemale lines of the typically monandrous species *Drosophila subobscura*, where males provide a nuptial gift. Heat stress significantly changed male gift presentation and female acceptance, impacting mating probability and reproductive output. Additionally, female remating probability rose significantly after copulating with a heat-stressed sterile male, with remating propensity varying across genotype. These findings suggest developmental heat stress affects multiple aspects of sex-specific reproduction, underscoring how climate change may influence sexual selection and mating system evolution.

**Authors:** Pembury Smith Matilda, Stockholm University, Sweden. Latkova Laura, Stockholm University, Sweden. Snook Rhonda, Stockholm University, Sweden.

**Key words:** sexual selection, mate choice, nuptial gift, temperature, mating system

**Email:** matilda.pembury.smith@zoologi.su.se

**Abstract number:** P9

**Title:** The stability of evolvability over 60 million years of Primate evolution

**Name:** Anna Penna

**University:** Lunds Universitet

**Abstract:**

The ability of evolving systems to change (evolvability) is deeply connected to the structure and amount of phenotypic variation transmitted from generation to generation (G-matrix). The G-matrix captures a population's immediate ability to evolve by selection and drift, making it a central parameter in quantitative genetics. Still, whether and how G evolves over prolonged evolutionary time scales is unclear. Here, we apply a comparative evolutionary quantitative genetics approach to investigate the stability of G in the primate radiation by comparing phenotypic covariance matrices (P-matrices). Based on 37 homologous 3D landmarks captured on the skull of 309 species (N = 9868 specimens), we estimated and compared P of 55 genera under a Bayesian framework. Our results show high levels of similarity, both in their variance structure and how they respond to selection. These findings suggest that even different shapes are governed by similar developmental systems.

**Authors:** Anna Penna (Lund University, Sweden) Fabio Machado (Oklahoma State University, USA)

**Key words:** G-matrix

**Email:** anna.penna@biol.lu.se

**Abstract number:** P10

**Title:** Context Dependent Male Mate Choice Copying in a Livebearing Fish

**Name:** Arezo Shamsgovara

**University:** Stockholm University

**Abstract:**

Mate choice drives evolution, often requiring costly assessment of ornamentation, which is indicative of mate quality. To avoid these costs, individuals can copy the mate choices of conspecifics. The phenomenon of mate choice copying is well established in females, yet its occurrence in males remains controversial. Male mate choice copying could offer benefits, such as identifying receptive females, but it also involves risks, such as increased sperm competition when choosing previously mated females. In this study, we explored male mate choice copying in a livebearing fish, pygmy halfbeak (*Dermogenys colletei*) using a dichotomous choice assay to see if a rival male's presence affects choices. Results suggest that male halfbeaks may copy the choice of conspecifics, but that this behaviour depends on social and ecological factors. This study sheds light on the context in which males mate choice copy, offering insights into the evolution of complex male mating behaviours.

**Authors:** Shamsgovara Arezo, Department of Zoology, Stockholm University, Sweden Winkler Lennart, Department of Zoology, Stockholm University, Sweden Wheatcroft David, Department of Zoology, Stockholm University, Sweden Kolm Niclas, Department of Zoology, Stockholm University, Sweden Fitzpatrick John L., Department of Zoology, Stockholm University, Sweden

**Key words:** Sexual selection, Mate choice, Mate Choice Copying, Male Mate Choice, Ornament

**Email:** arezosha2@gmail.com

**Abstract number:** P11

**Title:** Nuclear Cooperation as a Driver of Evolutionary Transitions in Fungi

**Name:** Mattias Siljestam

**University:** Stockholm University

**Abstract:**

The evolution of complex life has been shaped by transitions in individuality, where distinct entities combine to form higher-level organisms. In fungi, heterokaryosis—a condition where genetically distinct nuclei coexist in the same cytoplasm—provides an ideal system to study such transitions. My research focuses on *Neurospora tetrasperma*, a species that transitioned from homokaryosis to heterokaryosis approximately one million years ago. Using mathematical modelling within an evolutionary invasion analysis framework, I investigate how cooperation between different nuclear types drive this evolutionary shift. The study explore the conditions under which nuclear cooperation is favoured, providing insights into the mechanisms behind evolutionary transitions in individuality.

**Authors:** Siljestam, Mattias, Department of Ecology, Environment and Plant Sciences, Stockholm University, Sweden

Johannesson, Hanna, Department of Ecology, Environment and Plant Sciences, Stockholm University, Sweden

**Key words:** *Neurospora*, Adaptive Dynamics, Evolutionary Transitions, Multilevel selection

**Email:** mattias.siljestam@su.se

**Abstract number:** P12

**Title:** NA

**Name:** Rike Stelkens

**University:** Stockholm University

**Abstract:**

Accelerating climate change and extreme temperature events urge us to better understand the potential of populations to tolerate and adapt to new thermal challenges. Interspecific hybridization can facilitate adaptation to novel or extreme environments. However, predicting the long-term fitness effects of hybridization remains a major challenge in evolutionary and conservation biology. My lab uses experimental evolution with the powerful microbial model system budding yeast to track adaptive processes across hundreds of generations in real-time. We investigated the thermal adaptation dynamics of four species of budding yeast and their interspecific F2 hybrids, for 200 generations under extremely cold (5°C) and warm (31°C) conditions. We found significant variation in the evolutionary potential of species and hybrids. Cold-tolerant species showed larger fitness increases in warm temperature, whereas warm-tolerant species showed larger fitness gains in cold temperature. By far the largest fitness improvements occurred in hybrids, with some populations nearly quadrupling in fitness in the cold environment over the course of experimental evolution. Some hybrids exceeded both their parents in thermal adaptive potential. Reciprocal transplanting of evolved populations from the endpoint of evolution into opposite temperatures revealed that hybrids generally have greater resilience than their parents when challenged with sudden temperature shifts. Our results highlight that hybridization alters the fitness outcomes of long-term adaptation to extreme environments and may render populations more resilient to sudden environmental change, presenting both opportunities and challenges for conservation and sustainable agriculture.

**Authors:** Pinto, Javier; Stockholm University, Department of Zoology, Sweden  
Stelkens, Rike; Stockholm University, Department of Zoology, Sweden

**Key words:** climate change, experimental evolution, fitness, hybridization, yeast, thermal adaptation

**Email:** rike.stelkens@zoologi.su.se

**Abstract number:** P13

**Title:** Codon Usage Bias and Selection Patterns in Atlantic Herring

**Name:** Leyi Su

**University:** Uppsala University

**Abstract:**

Codon usage bias, which describes differences in the frequencies of occurrence of synonymous codons in coding sequences, is a prevalent finding in most organisms. The extent of codon usage bias is determined by many factors including gene expression level, GC content, tRNA abundance and effective population size. The Atlantic herring (*Clupea harengus*) is a model organism for exploring the genetic basis of ecological adaptation on account of its large effective population size. Our codon usage analyses demonstrate that the Atlantic herring shows stronger codon usage bias compared with Zebra fish and human. We then classified the synonymous mutations into three groups based on the direction of codon changes and found different selection patterns in the Site Frequency Spectra based on these three categories. However, the codon usage is largely affected by the GC3 content, so the drivers of observed stronger codon usage bias in Atlantic herring still requires more research.

**Authors:** Su Leyi, IMBIM, Uppsala University, Sweden; Wallberg Andreas, IMBIM, Uppsala University, Sweden; Pettersson Mats, IMBIM, Uppsala University, Sweden; Andersson Leif, IMBIM, Uppsala University, Sweden

**Key words:** Codon Usage Bias, GC content, Genome Evolution, Site Frequency Spectrum, Natural Selection

**Email:** leyi.su@imbim.uu.se

**Abstract number:** P14

**Title:** Natural and sexual selection in wild populations of tortoise beetles

**Name:** Maja Tarka

**University:** Lund University

**Abstract:**

Populations differ in their ability to adapt and evolve as a response to a changing environment. Understanding how and why these differences arise will be crucial for predicting their future. We use a novel study system for eco-evolutionary dynamics, the green tortoise beetle *Cassida viridis*, to understand how natural and sexual selection fluctuates between and within populations. By monitoring free ranging individual beetles in their natural habitat, we show that individuals are surprisingly sedentary, and this allows us to track them over their lives. This enables us to study how the environment, the host plant composition and phenotypic traits affect beetle survival and mating success. Our study provides valuable insights into the processes of fluctuating selection and adaptive strategies, elucidating the microevolutionary dynamics in wild insect populations.

**Authors:** Winslott, Erica, Dept. of Biology, Lund University, Sweden Botelho, Ana, Dept. of Biology, Lund University, Sweden Wendel, Erik, Dept. of Biology, Lund University, Sweden Lürig, Moritz, Florida Museum of Natural History, USA Karlsson Green, Kristina, Dept. of Plant Protection, SLU Alnarp, Sweden Tarka, Maja\*, Dept. of Biology, Lund University, Sweden

\*Presenting Author

**Key words:** microevolution, quantitative genetics, insects, survival and mating, phenotyping

**Email:** [maja.tarka@biol.lu.se](mailto:maja.tarka@biol.lu.se)

**Abstract number:** P15

**Title:** Population structure and isolation in different Swedish pollinator groups

**Name:** Isolde van Riemsdijk

**University:** Lund University

**Abstract:**

Insect numbers and diversity are in rapid decline, also within protected areas, endangering the ecosystem functions that insects provide and conflicting with societal commitment to protect biodiversity. Functional connectivity (the degree to which landscapes facilitate the movement of organisms, and the resulting fitness effect on populations) is crucial for population persistence, enabling three major evolutionary processes

**Authors:** van Riemsdijk, Isolde, Nolen, Zachery, Runemark, Anna et al. Lund University, Sweden

**Key words:** 1) re-colonization following stochastic extinctions

**Email:** [isolde.van\\_riemsdijk@biol.lu.se](mailto:isolde.van_riemsdijk@biol.lu.se)



**Abstract number:** P16

**Title:** Biodiversity Genomics Initiatives - sequencing life for the future of life

**Name:** Olga Vinnere Pettersson

**University:** SciLifeLab

**Abstract:**

Biodiversity genomics projects, in particular sequencing reference genomes, provide an open data resource to biologists worldwide. Reference genomes of eukaryotic life enable answering a plethora of biological questions. This resource enables ground-breaking research on, e.g. evolution of functions, conservation of endangered species, understanding the basic biology, etc. In this talk, I will present the international initiatives that Sweden is involved in, as well as give several examples of applications of reference genome sequencing for understanding of evolutionary processes.

**Authors:** Olga Vinnere Pettersson (UU-SciLifeLab, Sweden), Robert Waterhouse (SIB, Switzerland)

**Key words:** Genomics, open data, evolution, biodiversity

**Email:** olga.pettersson@scilifelab.uu.se

**Abstract number:** P17

**Title:** Inbreeding preference emerges in female guppies during more ecologically relevant mate encounters

**Name:** Amanda Viving

**University:** Stockholm University

**Abstract:**

Inbreeding is expected to reduce offspring fitness, known as inbreeding depression. Mate choice could help animals avoid inbreeding, but recent studies suggest that animals rarely use mate choice to avoid relatives, even when inbreeding is costly. Here, we examined the roles of sex, mating status, and social complexity on precopulatory inbreeding avoidance in guppies (*Poecilia reticulata*), a species experiencing inbreeding depression. Virgin and experienced male and female guppies were tested for their preference between related and unrelated partners in two experimental setups: (1) dichotomous choice and (2) free-swimming arena. In the dichotomous setup, neither sex preferred unrelated partners, but in the socially complex free-swimming arena females directed more sexual behaviors toward relatives. Mating status did not influence mate preference in either sex regardless of the experimental setup. These findings question the idea that mate preference shifts with sexual experience and underscores the importance of experimental design in researching mate choice.

**Authors:** Amanda Viving, Léa Daupagne, John Fitzpatrick. Stockholm University, Sweden

**Key words:** inbreeding avoidance, precopulatory mate choice, sex, mating status, social complexity

**Email:** amanda.viving@hotmail.com

**Abstract number:** P18

**Title:** Evolution and fine-scaled temperature variation in insect pest projections

**Name:** Loke von Schmalensee

**University:** Uppsala University

**Abstract:**

Climate change has caused changes in species' distributions, both in time and space. These shifts are projected to continue in the future, raising concerns about the spread of agricultural pests. Most climate projections are done by integrating phenotypic reaction norms with coarse climate projections, failing to incorporate evolutionary responses and important fine-scale temperature variation. Harnessing data from long-term experimental evolution under different thermal regimes in the insect pest, *Callosobruchus maculatus*, we project its future population growth and agricultural impact using 1-hour resolution thermal regimes, across 20 Californian sites. We use our framework to elucidate the relative contribution of evolution in different life-history traits to pest impact in a warmer world.

**Authors:** Loke von Schmalensee, Uppsala University, Sweden Elise Cazaux, Uppsala University, Sweden David Berger, Uppsala University, Sweden

**Key words:** Thermal adaptation, Climate change, Agricultural pests, Temporal variation

**Email:** loke.von.schmalensee@ebc.uu.se

**Abstract number:** P19

**Title:** eDNA in terrestrial animal monitoring

**Name:** Beilun Zhao

**University:** UPPSALA UNIVERSITY

**Abstract:**

Environmental DNA (eDNA) has become an invaluable tool for monitoring terrestrial animals, particularly for species that are challenging to detect through traditional methods. However, effective monitoring hinges on selecting appropriate sampling substrates and methods aligned with research objectives. The choice of substrate can influence species detection accuracy, affecting rates of false positives and negatives and capturing the spatiotemporal distribution of species. Our study investigates the detection biases introduced by substrates in assessing insect diversity, comparing eDNA substrates like anthill, sediment, soil, spiderweb, water, and decayed wood with traditional Malaise traps, commonly used to capture flying insects. Our analysis reveals that each substrate captures a unique taxonomic profile, with notable differences in diversity levels and species representation, underscoring the role of substrate choice in shaping biodiversity assessments. This study highlights the benefits of combining multiple substrates to reduce detection biases, enhance insect biodiversity assessments for improved ecological management and conservation strategies.

**Authors:** Beilun Zhao, Tobias Andermann Department of Organismal Biology Uppsala University Sweden

**Key words:** Insect Diversity, Environmental DNA (eDNA), Malaise Traps, Substrate Bias, Biodiversity Assessment

**Email:** beilun.zhao@ebc.uu.se

**Abstract number:** P20

**Title:** Understanding mitonuclear co-divergence through functional difference in *Ficedula* flycatchers

**Name:** Yishu Zhu

**University:** Uppsala University

**Abstract:**

Mitonuclear coevolution ensures metabolic function in organisms. Rapid-evolving mitochondrial genes is predicted to take the lead in this co-evolution and favour changes in nuclear alleles that either compensate for deleterious mitochondrial mutations (“nuclear compensation theory”) or reinforce favourable mitochondrial mutations that ensure adaptation to new environmental conditions (“nuclear reinforcement theory”). Here, I report co-divergence of mitonuclear OXPHOS genes encoding proteins building Complex I of the OXPHOS pathway in two flycatcher species as predicted by both hypotheses. However, high-resolution phenotypic data on cellular metabolic function reveal functional differences of Complex I activity that is more in line with the “nuclear reinforcement theory” and divergent climate adaptation in these wild populations. Such mechanism could contribute to speciation through the arise of mitonuclear incompatibility between lineages.

**Authors:** Yishu Zhu - Uppsala University, Sweden Eva van Der Heijden - University of Cambridge, UK Madeline Chase - Swiss Ornithological Institute, Switzerland Murielle Ålund - Uppsala University, Sweden S. Eryn McFarlane - University of Wyoming, US Antoine Stier - Université Lyon 1, France Tom van der Valk - Swedish Museum of Natural History, Sweden Carina Farah Mugal - UCB Lyon 1, France Anna Qvarnström - Uppsala University, Sweden

**Key words:** evolutionary biology, mitonuclear coevolution, *ficedula*, nuclear compensation

**Email:** yishu.zhu@ebc.uu.se

**Abstract number:** P21

**Title:** Do Sharks Respond to Eyespots?

**Name:** Madicken Åkerman

**University:** Stockholm University

**Abstract:**

Animal colourations and patterns exemplify evolutionary adaptations driven by predation to enhance prey survival. Eyespots, characterized by concentric rings of contrasting colours, have long been assumed to reduce predation by deceiving predators, and their anti-predator role has been demonstrated repeatedly. However, although well-studied in terrestrial environments, their impact on marine predators remains largely speculative. In this study, we demonstrate that eyespots elicit different responses in terrestrial predators (red junglefowl) and marine predators (brown-banded bamboo sharks and lemon sharks). Our findings reveal that terrestrial predators hesitate when confronted with eyespotted prey, whereas marine predators show accelerated attack responses. These contrasting behaviors suggest that eyespots function differently across environments, acting as deterrents in terrestrial settings but potentially attracting marine predators. Our study demonstrates that the function of eyespots varies across predator types and environments, emphasizing the need for further exploration in marine ecosystems where the function of eyespots is less understood.

**Authors:** Åkerman Madicken, Department of Zoology, Stockholm University, Sweden.

Juntorp Evelina, Department of Zoology, Stockholm University, Sweden.

Garnham Laura, Department of Physics, Chemistry and Biology, Linköping University, Sweden.

Løvlie Hanne, Department of Physics, Chemistry and Biology, Linköping University, Sweden.

Gotthard Karl, Department of Zoology, Stockholm University, Sweden.

Fitzpatrick John L., Department of Zoology, Stockholm University, Sweden.

**Key words:** conspicuous markings, detection-hypothesis, anti-predator colouration

**Email:** madicken.akerman@zoologi.su.se

**Abstract number:** P22

**Title:** The role of a structural variant for seasonal camouflage evolution in hares

**Name:** Huijuan Bi

**University:** Stockholm University

**Abstract:**

Close to 21 species of mammals and birds have convergently evolved the ability to seasonally change color, but it is unclear if the molecular mechanisms underlying this phenotype are also convergent across species. Previous genomic work in hares linked white to brown variation in winter coats to pigmentation gene ASIP. We revisited this genomic data and identified an indel in ASIP associated with the ability to change color in hares. We use comparative genomics to confirm the presence of this indel in the new PacBio reference-genome of the mountain hare. To explore the regulatory role of this mutation for the ability to change color, we annotate transcription factor binding sites (TFBS) in the indel region that are shared across color changing species. We identified several shared TFs in winter white hares, notably RBPJ, which is involved in maintaining proper hair pigmentation, and Prdm1, which plays a role in photoreceptor regulation.

**Authors:** Huijuan Bi, Department of Zoology, Stockholm University, Science for Life Laboratory, Sweden Mafalda S. Ferreira, Department of Zoology, Stockholm University, Science for Life Laboratory, Sweden

**Key words:** seasonal camouflage, ASIP, transcription factor

**Email:** huijuan.bi@zoologi.su.se

**Abstract number:** P23

**Title:** NA

**Name:** Omolola Afolabi

**University:** Federal College of Education Oyo State

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** firudngo@yahoo.com



**Abstract number:** P24

**Title:** On the origin of an insular hybrid butterfly lineage

**Name:** Jesper Boman

**University:** Uppsala Universitet

**Abstract:**

We report the discovery of a European hybrid butterfly lineage, a finding that can be considered surprising given the intense and long-term study of European butterflies. The lineage we describe is mainly inhabiting an island in the Baltic Sea and was previously designated as a subspecies (*horkei*) of one of the parental species (*Aricia artaxerxes*). By analysing whole-genome resequencing data, we determine that *horkei* originated by hybridization between the non-sister species *A. artaxerxes* and *A. agestis*. This hybridization event occurred approximately 54,000 years ago, predating the last glaciation of the current distribution range. *Horkei* must therefore have persisted long enough to be able to colonize its current range, despite that this area lies between the current distributions of the parental species. The hybrid origin, the maintenance of genomic integrity across times of dramatic climate change suggest that *horkei* could be in the process of hybrid speciation.

**Authors:** Jesper Boman<sup>1</sup>, Zachary J. Nolen<sup>2</sup>, Niclas Backström<sup>1</sup>

1: Evolutionary Biology Program, Department of Ecology and Genetics (IEG), Uppsala University, Norbyvägen 18D, SE-752 36 Uppsala, Sweden

2: Department of Biology, Lund University, Kontaktvägen 10, SE-223 62 Lund, Sweden

**Key words:** Biodiversity, Speciation, Hybridization, Hybrid speciation, Butterflies

**Email:** [jesper.boman@ebc.uu.se](mailto:jesper.boman@ebc.uu.se)

**Abstract number:** P25

**Title:** NA

**Name:** Andre Bourbonnais

**University:** Lund University

**Abstract:**

Insect herbivores exhibit remarkable diversity, with most species specializing on specific host plants. The intricate relationships between these herbivores and their host plants are strong drivers of insect diversification and speciation. Although the influence of microbial symbionts on these evolutionary processes is acknowledged, it remains unclear whether symbionts act as leading or adaptive factors in host divergence when adapting to new ecological niches.

To investigate the evolutionary processes that act on the symbiont as its host adapts to a novel host plant, I assembled and analyzed the metagenome-assembled genome (MAG) of the unculturable symbiont, *Candidatus Stammerula tephritidis* (Stammerula), which is maternally transmitted by the two host races in *Tephritis conura* (Diptera:Tephritidae). The assembled MAG showed characteristic signs of genome reduction typically found in obligate symbionts. In this study, the MAG of Stammerula was annotated and compared across different populations.

**Authors:** Bourbonnais, André, Lund University, Sweden

**Key words:** metagenomics, population genetics, symbiont, ecological speciation

**Email:** andbou95@gmail.com

**Abstract number:** P26

**Title:** Long-term genetic consequences of admixture in a sand lizard population

**Name:** Seraina Emilia Bracamonte

**University:** Uppsala University

**Abstract:**

The sand lizard (*Lacerta agilis*) occurs in fragmented populations in southern Sweden, including small, isolated populations that are of conservation concern. These populations have low genetic diversity and show signs of inbreeding depression. The semi-natural population of Stora Keholmen was established approximately 20 years ago by interbreeding different Swedish sand lizard populations. Stora Keholmen appears to have recovered from the inbreeding depression of its main parental population and has comparatively high genetic diversity. Here, we use whole genome sequencing to compare genetic diversity of the main parental population, the inbred Asketunnan population, to the admixed Stora Keholmen population. We determine genomic diversity within and between both populations and identify regions of high diversity in the admixed population. Similarly, we determine if the inbred population has retained diversity in some genomic regions, which may indicate ongoing selection.

**Authors:** Bracamonte, Seraina E., Uppsala University, Sweden Lindsay, Wilow R., Gothenburg University, Sweden Wapstra Erik, University of Tasmania, Australia Olsson, Mats, Gothenburg University, Sweden Lillie, Mette, Uppsala University, Sweden

**Key words:** Comparative genomics, admixture, conservation

**Email:** seraina.bracamonte@ebc.uu.se

**Abstract number:** P27

**Title:** The Swedish Reference Genome Portal: A new service facilitating access and discovery of genome data studied in Sweden

**Name:** Daniel Brink

**University:** SciLifeLab Data Centre

**Abstract:**

The Swedish Reference Genome Portal (<https://genomes.scilifelab.se/>) is a new service aiming to facilitate access and discovery of genome data of non-model eukaryotic species studied in Sweden. For each species, the portal aggregates taxonomic data, genome assemblies and annotation tracks, metadata, and links to external resources in a single place. The genome data is displayed in an embedded JBrowse 2 genome browser, allowing for visual representation of a variety of data types, and sharing of sessions between users. With this service, we aim to make genomics data more accessible for all users regardless of their prior level of bioinformatics knowledge, and promote FAIR data sharing. The poster will introduce key features of the Genome Portal and illustrate how researchers affiliated to Swedish institutions can submit their data. We are always interested in new additions to the portal! If you are interested, please contact us at [dsn-eb@scilifelab.se](mailto:dsn-eb@scilifelab.se).

**Authors:** Brink, Daniel P., SciLifeLab Data Centre, Uppsala University, Sweden Crean, Rory, SciLifeLab Data Centre, Uppsala University, Sweden Fuentes-Pardo, Angela P. , SciLifeLab Data Centre, Uppsala University, Sweden Ågren, Quentin, SciLifeLab Data Centre, Swedish Museum of Natural History, Sweden Kochari, Arnold, SciLifeLab Data Centre, Uppsala University, Sweden Lantz, Henrik, National Bioinformatics Infrastructure Sweden (NBIS), Uppsala University, Sweden Kultima, Hanna, SciLifeLab Data Centre, Uppsala University, Sweden Rung, Johan, SciLifeLab Data Centre, Uppsala University, Sweden Persson, Bengt, National Bioinformatics Infrastructure Sweden (NBIS), Uppsala University, Sweden

**Key words:** genomics

**Email:** [daniel.brink@scilifelab.se](mailto:daniel.brink@scilifelab.se)

**Abstract number:** P28

**Title:** Temporally stable parallel color cline in the shell of a marine gastropod from four coastal inlets

**Name:** Juan Gefaell

**University:** Lund University

**Abstract:**

The evolutionary ecology of *L. saxatilis*' shell color polymorphism has been relatively little studied. Here I describe a repeated color cline in the populations of *L. saxatilis* from the Rías Baixas (NW Iberian Peninsula), a series of four coastal inlets that arose from immersion of river valleys around 3000 years ago. We provide direct evidence that this color cline has remained relatively stable during the last 46 years. I also show, based on an analysis of snails from 55 localities across the four Rías Baixas, that there is a strong statistical association between the most prevalent color morphs and a suit of environmental and shell morphological variables, pointing to strong ecological selective pressures maintaining the cline and trait covariation. As the populations of *L. saxatilis* from the Rias Baixas got isolated relatively early, this putative example of parallel evolution has a bearing on discussions about predictability and contingency in evolution.

**Authors:** Juan Gefaell (1, 2), Ramón Vigo (1), Juan Galindo (1), Emilio Rolán-Alvarez (1)

- (1) Centro de Investigación Mariña, Universidade de Vigo, Departamento de Bioquímica, Genética e Inmunología, 36310 Vigo, Spain
- (2) Department of Biology, Lund University, Lund, Sweden

**Key words:** *L. saxatilis*, evolutionary ecology, color polymorphism, gastropod shell, parallel evolution

**Email:** [juan\\_gefaellborras@hotmail.com](mailto:juan_gefaellborras@hotmail.com)

**Abstract number:** P29

**Title:** Synonymous site constraint in mammals supports the unwanted transcript hypothesis

**Name:** Matthew Christmas

**University:** Uppsala University

**Abstract:**

It is known that selection acts on synonymous sites in mammals, but the functions held by synonymous sites are largely unresolved. Recently, the unwanted transcript hypothesis (UTH) has been presented as a potential explanation for evolutionary constraint at synonymous sites in species with low effective population sizes, such as humans. Selection for higher GC content and against mutations that alter splicing in native transcripts is predicted to shape synonymous site content and protect against unwanted transcripts. Utilising the Zoonomia alignment of 240 placental mammal genomes and single-base resolution constraint scores, we interpret mammalian synonymous site constraint in this context. We show that 20.8% of four-fold degenerate (4d) sites are under significant constraint, with a strong GC bias. We see evidence for higher 4d constraint in species with lower effective population sizes and greater genome transposable element content, as predicted by the UTH. Genes enriched for synonymous site constraint are integral to organismal viability through their involvement in embryo development and transcriptional regulation. Overall, our results provide support for the UTH and we conclude that it is the most viable model for explaining selection on synonymous sites in mammals

**Authors:** Matthew J. Christmas<sup>1,2</sup>, Michael Dong<sup>1,2</sup>, Jennifer R. S. Meadows<sup>1,2</sup>, Sergey Kosyrev<sup>1,2</sup>, Kerstin Lindblad-Toh<sup>1,2,3</sup> 1. Department of Medical Biochemistry & Microbiology, Uppsala University, Uppsala, Sweden 2. SciLifeLab, Uppsala University, Uppsala, Sweden 3. Broad Institute of MIT & Harvard, Cambridge, MA, USA

**Key words:** Evolutionary constraint, synonymous sites, mammals, GC content

**Email:** matthew.christmas@imbim.uu.se

**Abstract number:** P30

**Title:** NA

**Name:** Matthew Dalling

**University:** Linkoping University

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** DallingM@cardiff.ac.uk

**Abstract number:** P31

**Title:** NA

**Name:** Sylvester Aigbekaen

**University:** Federal College of Education Oyo State

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** bobadex010@outlook.com



**Abstract number:** P32

**Title:** Habitat complexity alters the strength of sexual selection on brain size in male and female halfbeaks (*Dermogenys collettei*)

**Name:** Léa Daupagne

**University:** Stockholm University

**Abstract:**

Animals often reproduce in complex environments, which should generate selection for increased detectability in signaling traits. However, the extent to which sexual traits have evolved to overcome the challenges of interacting in complex environments remains understudied. We examined whether habitat complexity influences sexual selection in the Pygmy Halfbeak, *Dermogenys collettei*, a small livebearing freshwater fish. Using free-swimming arenas, we created low- and high-complexity environments and observed mating behaviors in mixed-sex groups of three males and three females. While the opportunity for sexual selection did not differ significantly between environments for either sex, we detected positive selection for both male and female brain size in open arenas, but not in complex habitats. Interestingly, no clear selection on male or female ornamentation was detected in either environment. Our findings suggest that habitat complexity may limit selection pressures on cognitive traits, such as brain size, without affecting visual ornamentation's role in mating success.

**Authors:** Léa Daupagne, Stockholm University, Sweden Alessandro Devigili, Stockholm University, Sweden and University of Padova, Italy Rebecca McNeil, Stockholm University, Sweden John Fitzpatrick, Stockholm University, Sweden

**Key words:** Sexual selection, habitat complexity, mate choice

**Email:** lea.daupagne@zoologi.su.se

**Abstract number:** P33

**Title:** NA

**Name:** Gabriel David

**University:** Uppsala universitet

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** gbldavid@gmail.com

**Abstract number:** P34

**Title:** Innovative helmet complexity is phylogenetically conserved

**Name:** Marisol De la Mora Curiel

**University:** UNAM/ENES Juriquilla

**Abstract:**

The genomic data from ultraconserved elements were used to determine the evolutionary relationships among species of membracids, representing taxonomic diversity within the subfamilies and utilizing fresh specimens. The question of interest is whether the complexity of the helmet has evolved multiple times, or if organisms with similar helmet shape have a close evolutionary relationship, suggesting that the shape may be derived from a common ancestor. Our results show concordance with evolutionary relationships in previously published works

**Authors:** Ali Milanca Aguilar Gutierrez, Universidad Autónoma del Estado de Hidalgo (UAEH), Mexico. Esteban Alexis Vera Ortiz, Universidad Nacional Autónoma de Mexico (UNAM), Mexico. Adilson Pinedo Escatel, Universidad Nacional Autónoma de Mexico (UNAM), Mexico. Daniel Piñero Dalmaú, Universidad Nacional Autónoma de Mexico (UNAM), Mexico. Marisol De la Mora Curiel, Universidad Nacional Autónoma de Mexico (UNAM), Mexico.

**Key words:** the clades have a bootstrap support of 100, and the morphogeometric data support the hypothesis that similar pronota are found among phylogenetically close organisms

**Email:** mari.dc@unam.mx

**Abstract number:** P35

**Title:** Evolution of sexual dimorphism in resource use

**Name:** Stephen De Lisle

**University:** Karlstad University

**Abstract:**

Sex differences in ecologically important traits are common in animals and plants, and prompted Darwin to first propose an ecological cause of sexual dimorphism. Yet a role for ecological resource competition in the evolution of sexual dimorphism has never been directly demonstrated and remains controversial. I used experimental evolution in *Drosophila melanogaster* to test the hypothesis that resource competition can drive the evolution of sex differences in diet. Over just a handful of generations, offspring from flies evolved in low-resource, high-competition environments show elevated sexual dimorphism in diet preference compared to both the ancestor and populations evolved on high-resource availability. Expanding this design to include a manipulation of sex ratio indicates that over longer timespans diet evolves in predictable ways in response to operational sex ratio. These results indicate an important role for both ecology and mating system in driving the evolution of sexual dimorphism in ecologically important traits

**Authors:** Stephen P. De Lisle

**Key words:** Sexual dimorphism, evolutionary ecology, quantitative genetics

**Email:** [stephen.de.lisle@kau.se](mailto:stephen.de.lisle@kau.se)

**Abstract number:** P36

**Title:** Exploratory behavior in the hybrid Italian sparrow (*Passer italiae*) and its parental species

**Name:** Samantha Duhé Jones

**University:** Lund University

**Abstract:**

To address how behavioral traits evolve in hybrid lineages, we use the homoploid hybrid Italian sparrow (*Passer italiae*), whose parental species are the commensal house sparrow (*P. domesticus*) and non-commensal Spanish sparrow (*P. hispaniolensis*). We use a novel environment test to investigate whether exploration in hybrids is intermediate, mosaic, or transgressive using lineages of Italian sparrow from Crete, Corsica, and Sicily, differing in parental genetic contribution, as well as the two parental species and male F1 hybrids. We recorded time spent on features, latency to first feature, number of unique features visited, total visits, and two principal components derived therefrom. Across measurements, the most important predictor is the interaction of population with sex. House sparrows and Italian sparrow females are more active than Spanish sparrows and Corsican and Sicilian males, while Cretan males are transgressively active. In conclusion, both commensal Italian sparrows and F1 males generally have intermediate exploratory behavior.

**Authors:** Duhé Jones, Samantha, Lund University, Sweden Sokolovskis, Kristaps, Lund University, Sweden Steward, Rachel, Lund University, Sweden Hohne, Veronica, Netherlands Institute of Ecology, Netherlands Lawrence, Alex, Lund University, Sweden Wessman, Petronella, Lund University, Sweden Tunström, Kalle, Lund University, Sweden Rowe, Melissa, Netherlands Institute of Ecology, Netherlands Ravinet, Mark, University of Oslo, Norway Runemark, Anna, Lund University, Sweden

**Key words:** hybridization, behavior, commensalism, intermediacy, transgression

**Email:** sa1327du-s@student.lu.se

**Abstract number:** P37

**Title:** Discovering genotype-phenotype relationships with machine learning and the Visual Physiology Opsin Database (VPOD)

**Name:** Seth Frazer

**University:** University of California Santa Barbara

**Abstract:**

Predicting gene function from genetic variation is crucial for fields like bioengineering and evolutionary biology. Opsins, light-sensitive proteins affecting phenotypes like color vision, are a well-studied model gene family. However, existing opsin data is scattered and inaccessible, hindering systematic analysis.

We introduce the Visual Physiology Opsin Database (VPOD), compiling 1211 unique opsin genotypes and corresponding light absorption phenotypes (lambda-max) from 111 publications. Using VPOD and machine learning, we demonstrate accurate lambda-max prediction, accounting for non-additive mutation effects, and identification of critical amino acid sites.

This ability to predict gene function from sequence alone allows exploration of molecular evolution, ecological connections, and protein design. VPOD and our models provide a foundation for future research on gene families with quantifiable phenotypes.

**Authors:** Seth A. Frazer<sup>1</sup>, Mahdi Baghbanzadeh<sup>2</sup>, Ali Rahnavard<sup>2</sup>, Keith A. Crandall<sup>2,3</sup>, Todd H. Oakley<sup>1</sup>

**Affiliations:** 1 - Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, California 93106 2 - Computational Biology Institute, Department of Biostatistics and Bioinformatics, Milken Institute School of Public Health, The George Washington University, Washington, DC 20052 3 - Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC 20012

**Key words:** Machine learning

**Email:** safrazer7722@gmail.com

**Abstract number:** P38

**Title:** Can co-flowering species influence on each other's evolution? Insights from food-deceptive orchids in Öland

**Name:** Yedra García

**University:** Lund University

**Abstract:**

Interactions between co-flowering species for shared pollinators are common and can influence plants' reproductive success and ultimately natural selection acting on floral traits. However, studies testing for pollinator-mediated reproductive interactions in local communities are rare mainly due to lack of appropriate statistical methods. Here, we applied recent statistical models within the Hierarchical Modelling of Species Communities (HMSC) framework to test for pollinator-mediated interactions among three co-flowering species and estimate phenotypic selection on floral traits simultaneously on the three species. To do that, we recorded phenotypic traits, flower abundance and reproductive fitness of three food-deceptive co-flowering orchid species across 75 1m<sup>2</sup> patches with varying species composition. Our study showed species-species associations influencing the reproductive fitness of plant species at local communities and yield new insights on the biotic factors influencing the success of food-deceptive pollination systems. Moreover, our work provides a promising approach towards understanding natural selection at the community level.

**Authors:** García Yedra, Lund University, Department of Biology, Sweden Friberg Magne, Lund University, Department of Biology, Sweden Edrén Mikael, Lund University, Department of Biology, Sweden Opedal Øystein H., Lund University, Department of Biology, Sweden

**Key words:** reproductive interactions, pollinators, natural selection, orchids, community

**Email:** yedra.garcia\_garcia@biol.lu.se

**Abstract number:** P39

**Title:** Taming Evolution

**Name:** Johanna Gjoen

**University:** Linkoping University

**Abstract:**

Gjoen, Johanna, Linkoping University, Sweden Jensen, Per, Linkoping University, Sweden

**Authors:** The impact of selection and domestication on behaviour in Red Junglefowl

**Key words:** We investigated long-term selection effects on tameness in Red Junglefowl (*Gallus gallus*), at the 13th generation, using 40 birds: 20 from each line, with equal male-female distribution in the high-fear line, and 12 males and 8 females in the low-fear line. Previous studies have shown significant differences in cerebellum size and neuron density between Red junglefowl selected for high or low-fear of humans. Hence, we focused on traits associated with the cerebellum, including motor coordination, balance, as well as social navigation, to assess selection impacts. Behavioural tests (obstacle course, rotarod, and Y-maze) were used to explore the correlation between tameness levels and cerebellar function. Preliminary observations indicate potential motor and social trait differences between selection lines, offering insights into cerebellar adaptability under selective pressures. Complete results will provide more insight into the relationship between cerebellum-linked behaviours and selection, as well as cerebellar adaptation in evolutionary processes.

**Email:** Johanna.gjoen@liu.se



**Abstract number:** P40

**Title:** Meta-analytical evidence for frequency-dependent selection across the tree of life

**Name:** Miguel Gómez

**University:** Karlstad University

**Abstract:**

Explaining the maintenance of genetic variation in fitness-related traits within populations is a fundamental challenge in ecology and evolutionary biology. Frequency-dependent selection (FDS) is one mechanism that can maintain such variation, especially when selection favours rare variants (negative FDS). However, our general knowledge about the occurrence of FDS, its strength and direction remain fragmented, limiting general inferences about this important evolutionary process. We systematically reviewed the published literature on FDS and assembled a database of 747 effect sizes from 101 studies to analyse the occurrence, strength, and direction of FDS, and the factors that could explain heterogeneity in FDS. Using a meta-analysis, we found that overall, FDS is more commonly negative, although not significantly when accounting for phylogeny. An analysis of absolute values of effect sizes, however, revealed the widespread occurrence of modest FDS. However, negative FDS was only significant in laboratory experiments and non-significant in mesocosms and field-based studies. Moreover, negative FDS was stronger in studies measuring fecundity and involving resource competition over studies using other fitness components or focused on other ecological interactions. Our study unveils key general patterns of FDS and points in future promising research directions that can help us understand a long-standing fundamental problem in evolutionary biology and its consequences for demography and ecological dynamics.

**Authors:** Gómez-LLano, Miguel. Karlstad University, Sweden Bassar, Ronald. Auburn University, USA Svensson, Erik. Lund University, Sweden Tye, Simon. University of Arkansas, USA Siepielski, Adam. University of Arkansas, USA

**Key words:** frequency dependent selection

**Email:** miguel.gomez@kau.se

**Abstract number:** P41

**Title:** Unravelling the Cost of Pigmentation: Gene Knockouts and Their Effects in *Pieris brassicae*

**Name:** Eleine Gómez Sundberg

**University:** Stockholm University

**Abstract:**

Butterflies are known for their vibrant colouration, but the costs of producing these colours is not well understood. *Pieris brassicae* use pteridine for their white pigments, which are stored in pigment granules known as pterinosomes. Transport of pteridine precursors is mediated by an ABC transporter complex formed through the heterodimerization of white and brown. In lepidoptera, the *ok* gene, a paralog for brown also heterodimerize with white. To explore the functional role of these genes, we used CRISPR/Cas9 to knockout white, brown, and *ok* in *P. brassicae* embryos. Mutant phenotypes resulted in various developmental and pigmentation defects (translucent larval epidermis, pupal colour disruption). Adult butterflies displayed suppressed wing development, and absence of wing pigmentation, along with distinct eye colour changes. These findings reveal the relative roles of the genetic mechanisms behind butterfly pigmentation and ultimate shed light on the cost of producing colour.

**Authors:** Gómez-Sundberg, Eleine R., Department of Zoology, Stockholm University, Sweden Medina-Jiménez, Brenda I., Department of Zoology, Stockholm University, Sweden Wheat, Christopher W., Department of Zoology, Stockholm University, Sweden

**Key words:** Pteridine Transport, Butterfly coloration, ABC transport complex, CRISPR/Cas9

**Email:** Eleine.gs@gmail.com

**Abstract number:** P42

**Title:** Lens canariensis - Two millennia of lentil cultivation on the Canary Islands

**Name:** Jenny Hagenblad

**University:** Linköping University

**Abstract:**

During more than a millennium, agriculture in the Canary Islands relied on a limited number of crop species. During this time the islands were isolated from each other and the mainland making each island an independent evolutionary replicate of crop plant - human - nature interrelations. The European discovery and conquest of the islands in the 15th century turned a Neolithic society into a globalisation hub between the old and new world with dramatic consequences for the human population and for agriculture.

Through genetic analyses of present-day genebank accessions, combined with aDNA analyses of specimens more than a millennium old, the evolution of the Canarian lentil has been investigated. The results show the consequences of isolation and globalisation as well as the effects of both climate and human society and hint to a role of Canarian lentils in the cultivation and consumption of lentils far beyond the Canarian archipelago.

**Authors:** Hagenblad, Jenny, Linköping University, Sweden Morales, Jacob, University of Las Palmas de Gran Canaria, Spain Fregel, Rosa, University of La Laguna, Spain Leino Matti W., Stockholm University, Sweden Rodríguez-Rodríguez, Amelia C., University of Las Palmas de Gran Canaria, Spain Santana Cabrera, Jonathan, University of Las Palmas de Gran Canaria, Spain

**Key words:** aDNA, crop evolution, agrarian history, plant - human interrelations

**Email:** Jenny.Hagenblad@liu.se

**Abstract number:** P43

**Title:** Cost of spontaneous dispersal is not sufficient to drive adaptive evolution in a spatially structured habitat

**Name:** Subhasish Halder

**University:** Linköping University

**Abstract:**

How populations adapt to the ecological challenges related to spatial-structuring— a common attribute of most naturally occurring populations, is poorly understood. Previously, using laboratory adapted populations of *Drosophila melanogaster* I have shown fitness cost of living under spatial-structuring owing to sex-biased spontaneous dispersal. Here, I took an experimental evolution approach to investigate the evolvability of spontaneous dispersal. Following 30+ generations of maintenance in a structured habitat with inter-patch movement, I found experimental and control populations do not differ in dispersal related traits. Females from experimental populations continued to pay reproductive and survival costs when subjected to experimental habitat structuring, indicating no sign of adaptation to population structuring. I argue that adaptation to habitat structuring depends on the intricate details of an ecology and discuss the problems in the detection of heritability in spontaneous dispersal including non-transitivity in dispersal syndrome, and the caveats and confounds of my experimental paradigm.

**Authors:** Subhasish Halder , Anupam Barman, Bodhisatta Nandy\*

Indian Institute of Science Education and Research Berhampur, Transit Campus, Govt. ITI Building, NH 59, Engineering School Junction, Ganjam, Berhampur 760 010, Odisha, India.

**Key words:** Population spatial-structuring, Sex-biased dispersal, Experimental evolution

**Email:** subhasish.halder@liu.se

**Abstract number:** P44

**Title:** Adapting fast and slow: The repeatability and molecular basis of plastic and genetic responses to temperature extremes

**Name:** Alex Hart

**University:** Uppsala University

**Abstract:**

Theory predicts that phenotypic plasticity can play a pivotal role in evolution by impeding or facilitating genetic adaptation. Thus, to predict how populations will respond to climate change, we need to understand how plastic and genetic responses interact during thermal adaptation.

We used experimental evolution in seed beetles to unravel plastic and genetic responses to temperature stress. Replicate lines were generated from three geographic origins, and then adapted to cold, ancestral, or hot temperature for ~100 generations. Gene expression was then measured at the three temperatures.

In line with previous findings, genetic changes in expression often counteracted immediate plastic responses, suggesting that genetic compensation is a major mode of thermal adaptation across the transcriptome. However, this pattern was much stronger and more repeatable across genetic backgrounds for cold-adaptation compared to heat-adaptation. We harness pool-sequencing and phenotypic data from the same lines to explore the underlying mechanisms behind these qualitative differences.

**Authors:** Hart, Alex: Uppsala University, Sweden Rego, Alex: Uppsala University, Sweden Stelkens, Rike: Stockholm University, Sweden Berger, David: Uppsala University, Sweden

**Key words:** genomics, climate change, thermal adaptation, genetic plasticity

**Email:** alex.hart@ebc.uu.se

**Abstract number:** P45

**Title:** Gone with the wind: exploring a vanished African rock doves hybrid zone in the Sahara Desert

**Name:** Germán Hernández Alonso

**University:** Uppsala University

**Abstract:**

A previous genomic study of rock doves (*Columba livia*) identified a genetically distinct population in West Africa, possibly representing a new species. This divergence is hypothesized to have arisen through cycles of allopatry during the dry and wet Sahara periods. Based on the Refugia Theory and observed admixture patterns, it was proposed that a hybrid zone existed in the Sahara during its last green period. This project aims to test the existence and location of this extinct hybrid zone by analyzing whole-genome sequences from six historical rock doves from previously unsampled populations in the Central Sahara and West Africa, in addition to existing genomic data. The findings confirm the presence of the hybrid zone, with a likely location in the mountainous and hilly regions of northwest Africa. Additionally, the results reveal a strong population structure and confirm the genetic divergence of West African rock doves.

**Authors:** Hernández-Alonso Germán, Uppsala University, Sweden. Feizabadi Farahani Motahare, Stockholm University, Sweden. van Grouw Hein, Natural History Museum, UK. Günther Torsten, Uppsala University, Sweden.

**Key words:** Population Genomics, *Columba livia*, Speciation, Refugia theory

**Email:** german.hernandez-alonso@uu.se

**Abstract number:** P46

**Title:** NA

**Name:** Amparo Herrera-Dueñas

**University:** Lund University

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** amparo.herrera\_duenas@biol.lu.se

**Abstract number:** P47

**Title:** Genetic Mechanisms of Host Shifting in *Acrocercops transecta*: A Linkage Map Analysis Using Gras-Di

**Name:** Bin Hirota

**University:** Kyoto Prefectural University

**Abstract:**

Host shifts in phytophagous insects often leads to genetic divergence between populations and adaptation to a novel host plant can promote speciation. A leaf-mining moth, *Acrocercops transecta*, consists of two host races that are associated with either Juglandaceae or *Lyonia ovalifolia* (Ericaceae). To understand the genetic background of shifting to *Lyonia*, backcross hybrids were propagated between the two host races. Genotyping by Random Amplicon Sequencing-Direct (Gras-Di) was used to construct a linkage map and to identify genetic regions responsible for oviposition preference and larval performance. We will show the genomic architecture of host shifting between distantly related host plants and discuss its consequences for ecological speciation in *A. transecta*.

**Authors:** Bin Hirota, Issei Ohshima (Kyoto Prefectural University, Japan)

**Key words:** host race, speciation, phytophagous insects, larval performance, oviposition preference

**Email:** [hirotabin@gmail.com](mailto:hirotabin@gmail.com)



**Abstract number:** P48

**Title:** Skull bones of contrasting developmental origins influence both micro- and macroevolution in lacertid lizards

**Name:** QUENTIN HORTA-LACUEVA

**University:** Lund University

**Abstract:**

Vertebrates are remarkable for their diversity of skull shapes, the evolvability of which might be determined by the specific variational properties of bones conferred by two types of embryonic tissues (the neural crest vs. the mesoderm). We tested this hypothesis by studying skull shape in the highly diverse family of Lacertid lizards, using X-ray microtomography-based (CT) 3D morphometric analyses. At the microevolutionary scale, we quantified variation in skull shape across populations of European wall lizards (*Podarcis muralis*) known to exhibit extreme morphological variations. At the macroevolutionary scale, we compare skull shape across more than 150 lacertid species. We quantified the contribution of the main skull bones to patterns of variation with regards to their neural crest or mesoderm origin. Family-level disparity patterns concordant with the variability observed in *P. muralis* would reveal if and how the neural crest biology underlies the possible integration of micro- and macroevolutionary processes.

**Authors:** Quentin Horta-Lacueva, Lund University, Sweden Juliane Vehof, Leibniz Institute for the Analysis of Biodiversity change, Museum Koenig Bonn, Germany Morris Flecks, Leibniz Institute for the Analysis of Biodiversity change, Museum Koenig Bonn, Germany Tobias Uller, Lund University, Sweden Nathalie Feiner, Lund University, Sweden - Max Planck Institute for Evolutionary Biology, Germany

**Key words:** skull morphology, developmental biases, microevolution, macroevolution, geometric morphometrics

**Email:** quentin.horta-lacueva@biol.lu.se

**Abstract number:** P49

**Title:** NA

**Name:** Arild Husby

**University:** Uppsala University

**Abstract:**

Understanding how environmental signals are sensed by organisms and how such signals lead to regulatory variation that induce changes in phenotypes are important if we are to understand how organisms adapt to changing environments. In many insects a particular case of this is wing polyphenism, the environmentally induced development of small or large wings. This is observed in several different waterstriders and we have combined tissue specific transcriptomic analyses with functional genomic tools to demonstrate that the Fat/Hippo pathway control this wing polyphenism in *Gerris buenoi*. How this occur is however unknown and here we examine the potential role of alternative splicing as well as hormonal assays to try to identify the upstream regulator of Fat/Hippo. Our results implicate the involvement of a new GRN in wing polyphenism and demonstrate the diversity of regulatory mechanism involved, despite a shared phenotypic basis across hemimetabolous insects for wing polyphenism.

**Authors:** Gudmunds, Erik, UU, Sweden Palahi I Torres, Aleix, UU, Sweden Khila, Abderrahman, UU/IGFL, Sweden/France Husby, Arild, UU, Sweden

**Key words:** phenotypic plasticity, gene regulation, developmental plasticity, alternative splicing

**Email:** Arild.husby@ebc.uu.se

**Abstract number:** P50

**Title:** Conservation genomics of wall lizards (*Podarcis muralis*) in the UK

**Name:** Santiago Jose Avila Quintero

**University:** Lund University

**Abstract:**

Small populations experiencing inbreeding depression may initiate a purging process, where natural selection gradually eliminates strongly deleterious mutations. Introduced populations frequently exhibit a lag phase preceding their expansion, which may involve purging. Here we studied wall lizards (*Podarcis muralis*) in the UK, where multiple introductions from France and Italy have given rise to isolated, small populations. These offer a unique opportunity to study their population genomics and test for purging of deleterious mutations using independent replicates. We will present our initial results on the genetic structure of these populations, using analyses based on whole-genome data from both introduced and native populations. These analyses will help identify the precise origins of the introduced populations. Additionally, we are assessing genomic diversity and investigating signs of inbreeding across these populations. Our ongoing exploration includes examining deleterious mutations and potential purging effects, providing unique insights into the genetic status of these populations.

**Authors:** Avila-Quintero Santiago, Lund University, Sweden Uller Tobias, Lund University, Sweden Feiner Nathalie, Max Planck Institute for Evolutionary Biology, Germany

**Key words:** Conservation genomics, evolutionary genomics, introduced species, Wall lizards, purging.

**Email:** sanjosaviqui@gmail.com

**Abstract number:** P51

**Title:** Metagenomic reconstruction of the mammoth steppe flora from 48,000 BP based on the steppe bison intestinal content

**Name:** Chenyu Jin

**University:** Centre for Palaeogenetics

**Abstract:**

During the last Pleistocene cold stage (115,000 - 11,700 years ago), the non-glaciated part of Arctic Eurasia exhibited a distinctive plant composition consisting of steppe and xerophilous arctic plants adapted to a dry and cold climate. This vegetation formed the Pleistocene mammoth steppe, which provided a habitat for large herbivores like mammoths (*Mammuthus primigenius*) and steppe bison (*Bison priscus*). In this study, we analyzed a plant paleo-community of the Arctic mammoth steppe through DNA sequencing of the well-preserved intestinal content in a bison ice mummy, known as the Alazeya steppe bison, which was discovered in the Kolyma Lowlands in northeast Russia in 2009 and dated to >48,000 14C years before present. We leveraged the power of plant genome-wide and skim genome accessions to maximize the retrieval of the sequence reads belonging to plant genera. Macrofossil analysis of this sample revealed a plant composition resembling a saline meadow. In line with macrofossil evidence, we identified alkali grass (*Puccinellia* sp.) indicating high salinity, and Larch (*Larix* sp.) in the DNA data of the steppe bison sample. The Larch (*Larix*) and the range of most likely species of willow (*Salix*) are indicative of cold temperatures and paludification. We also confirmed the presence of plant genera typically associated with arid soil conditions such as *Artemisia* and *Koeleria*, while genera *Phippsia*, *Alopecurus*, and *Festuca* could suggest high soil water content, at least seasonally. Some of these taxa were not previously identified in this material by macrofossil analyses. Hence, the use of metagenomic approaches combined with comprehensive reference databases allows us to obtain a more detailed view of the mammoth steppe vegetation and provides extended interpretations of soil conditions.

**Authors:** Chenyu Jin, Adrian Forsythe, Miklós Bálint, Frank Kienast, Pavel A. Nikolskiy, Rafał Kowalczyk, Katerina Guschanski, Tom van der Valk

**Key words:** shotgun sequencing, metagenomic, Middle Pleistocene

**Email:** chenyu.jin@nrm.se

**Abstract number:** P52

**Title:** Long-term use of Concept Inventory of Natural Selection on Swedish University students shed light on difficult concepts and identifies improved English skills over the years

**Name:** Anna M Johansson

**University:** Swedish University of Agricultural Sciences

**Abstract:**

A concept inventory is a set of multiple-choice questions, where the wrong answers correspond to common misconceptions. We translated the Concept Inventory of Natural Selection (CINS) to Swedish and used have used the English and Swedish versions since 2012. The concept "Change in a population" has consistently been most difficult for the students with the misconception "Learned behaviors are inherited" getting the most answers alongside "Mutations occur to meet the needs of the population". Interestingly, the same question asked about different species indicates that choice of species in a question can have an impact on how students perceive a concept. Overall, there are differences between students that gets the questions in Swedish or English. During the first years it took the students longer time to answer questions posed in English than in Swedish. In recent years the difference in answering time has disappeared, indicating an improved ability to read English.

**Authors:** Johansson Anna M 1 and Hagenblad Jenny 2 1Swedish University of Agricultural Sciences 2Linköping University

**Key words:** evolution, education, language

**Email:** anna.johansson@slu.se

**Abstract number:** P53

**Title:** Avian cognition: the influence of sleep

**Name:** Robin Johnsson

**University:** Franklin & Marshall College

**Abstract:**

Cognition, encompassing the processes through which animals acquire, store, and use information, is vital for survival across taxa. Notably, cognitive functions vary both between and within species, yet the underlying mechanisms remain largely unexplored. This presentation focuses on sleep, a critical but often overlooked neurophysiological phenomenon. Remarkably, despite a separation of 300 million years of evolution, both mammals and birds have independently evolved two unequivocal sleep states: non-rapid eye movement (NREM) sleep and REM sleep. Research indicates that these sleep states enhance cognitive processes in both groups, particularly in relation to learning and memory consolidation. During wakefulness, new memory traces are acquired and encoded in the brain, and subsequent sleep stabilizes and enhances newly formed memories. Conversely, sleep loss can hinder these neurological processes, resulting in impaired cognitive functions required for ideal behavioural performance. In this presentation, I will summarize the current research on sleep-dependent cognition in birds.

**Authors:** Johnsson, Robin, Franklin & Marshall College, USA

**Key words:** cognition, sleep, birds

**Email:** johnsson.rd@gmail.com

**Abstract number:** P54

**Title:** NA

**Name:** Ibraheem Kunle Azeez

**University:** Federal College of Education Oyo State

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** [info@firundfoundation.org](mailto:info@firundfoundation.org)

**Abstract number:** P55

**Title:** The role of chromosomal inversions in adaptation to salinity stress in Atlantic cod

**Name:** Rebecca Krohman

**University:** University of New Brunswick

**Abstract:**

Recombination is suppressed within inversions allowing these chromosomal regions to evolve separately and causing them to spread quickly through populations if they allow an individual to adapt better to local conditions. Individuals who contain an inversion with locally adapted alleles will be more likely to survive in a population exposed to environmental stressors. Atlantic cod is an ecologically, culturally, and economically important species and classified as endangered due to previous overfishing. Cod contains multiple inversions throughout its genome. Using common garden experiments, I will identify when mortality occurs and how inversion genotype frequencies change when the early life of cod are exposed to salinity stress. This study will directly test which genomic regions influence the survival of early life stages of cod in specific environmental conditions, allowing for strong inferences regarding which genotypes are adaptive. My results will help inform management and conservation strategies to increase cod population size.

**Authors:** Rebecca Krohman, University of New Brunswick, Canada

Rebekah Oomen, University of New Brunswick, Canada; University of Gothenburg, Sweden; University of Oslo, Norway

**Key words:** Atlantic cod, Chromosomal inversion, salinity stress, common garden experiment

**Email:** rebecca.Krohman@unb.ca



**Abstract number:** P56

**Title:** The frequency, drivers and consequences of introgression among Swiss ants revealed by large-scale genotyping

**Name:** Guillaume Lavanchy

**University:** Lund University

**Abstract:**

Hybridization and the resulting introgression are increasingly recognized as major forces driving species and genome evolution, but estimating its frequency in nature is challenging. Here, we genotyped over 4100 individuals belonging to 40 species from seven ant genera using ddRAD sequencing to screen for signatures of admixture, indicative of contemporary introgression. We found that the proportion of species engaging in introgression is much higher than previous estimates, but the amount of introgression is very low in most cases. This suggests that many species introgress a little, contrasting with previous findings of abundant hybridization between only a few species. We found support for the prediction that introgression is more frequent between species with overlapping mating phenology and low genetic divergence, indicating that both behavioural barriers (pre-mating) and incompatibilities (post-meiotic) play important roles in shaping introgression in these ants. Introgression contributed to overall genetic diversity, but moderately in most cases.

**Authors:** Guillaume Lavanchy, Biology Department, Lund University, and Department of Ecology and Evolution, University of Lausanne, Switzerland Ludovic Ruedi, Department of Ecology and Evolution, University of Lausanne, Switzerland Kristine Jecha, Department of Ecology and Evolution, University of Lausanne, Switzerland Jérôme Goudet, Department of Ecology and Evolution, University of Lausanne, Switzerland Tanja Schwander, Department of Ecology and Evolution, University of Lausanne, Switzerland

**Key words:** introgression, hybridization, evolutionary genomics

**Email:** guillaume.lavanchy@biol.lu.se

**Abstract number:** P57

**Title:** Investigating the segregation patterns of gCNVs

**Name:** Freja Lindstedt

**University:** Uppsala University

**Abstract:**

Gene copy number variations are pervasive in eukaryotic genomes, but understudied. Open evolutionary questions include gCNV duplication rates, segregation patterns and impact on recombination. Previously we identified gCNVs across 25 Brassicaceae species. Using this, I have designed crosses to investigate the evolution of target gCNVs in *Capsella rubella*.

**Authors:** Lindstedt Freja, Uppsala University / SciLifeLab, Sweden Milesi Pascal, Uppsala University / SciLifeLab, Sweden

**Key words:** gCNVs, evolution, Brassicaceae

**Email:** freja.lindstedt@ebc.uu.se

**Abstract number:** P58

**Title:** NA

**Name:** Adetayo Samuel Babalola

**University:** Federal College of Education Oyo State

**Abstract:**

NA

**Authors:** NA

**Key words:** NA

**Email:** bobadex010@gmail.com

**Abstract number:** P59

**Title:** Flower allometry and its relationship with pollinators along an altitudinal gradient

**Name:** Sissi Lozada Gobillard

**University:** Lund university

**Abstract:**

Flower traits have been shown to be related to breeding systems (outcrossing vs. selfing), pollinators and can vary along environmental gradients. Environmental variation along altitudinal gradients is commonly associated with changes in the diversity of plants and animals, making them ideal systems to study variation in flower traits and its relationship with pollinators. We measured flower traits of 20 plant species and observed their pollinators (flower visitors that touched reproductive parts of the flower) along an altitudinal gradient (1000-4500m) in the tropical Andes of Bolivia. Flower traits included flower size, entrance diameter, flower length and anther-stigma distance. In particular, we compared allometries of stigma-anther height along the gradient. In this study we test the hypothesis that anther-stigma correlations is stronger at higher altitudes, assuming that at higher altitudes the conditions are harder, pollination is less reliable and therefore selfing plants species would be more common. Additionally, we tested whether allometric slopes for fit traits are shallower than size traits, and how they vary among pollinators.

**Authors:** Sissi Lozads Gobillard 1, 2 Oystein Opedal 1 1 Biodiversity and Evolution, Department of Biology Lund University 2 instituto de Ecologia, Universidad Mayor de San Andres La Paz Bolivia

**Key words:** Flower traits, flower evolution, altitudinal gradient, tropical Andes, Bolivia

**Email:** [sissi.lozada\\_gobillard@biol.lu.se](mailto:sissi.lozada_gobillard@biol.lu.se)

**Abstract number:** P60

**Title:** Single-cell sequencing suggests a conserved function of Hedgehog-signalling in spider eye development

**Name:** Brenda Irene Medina Jiménez

**University:** Stockholm University

**Abstract:**

Spiders evolved different types of eyes, commonly called primary and secondary eyes. Arthropod eye development is best understood in the vinegar fly *Drosophila melanogaster*, which evolved three types of eyes, the larval eyes, ocelli and compound eyes. The gene regulatory networks that underlie their development exhibit a conserved core network that attest of similarities between these eyes. Using members of this core network as candidate genes, studies have revealed crucial differences such as the apparent lack of some key factors in some groups of arthropods, including spiders. By analysing cell transcriptomes from spider embryos undergoing organogenesis, we found that Hedgehog (Hh)-signalling seems to be involved in eye development in the spider *Parasteatoda tepidariorum* and demonstrated that this involvement is conserved in spiders. Additionally, we identified genes that are expressed in the developing eyes of spiders, but that have not been studied in this context before.

**Authors:** Medina-Jiménez Brenda I., Stockholm University, Sweden Budd Graham E., Uppsala University, Sweden Pechmann Matthias, University of Cologne, Germany Posnien Nico, University of Göttingen, Germany Janssen Ralf, Uppsala University, Sweden

**Key words:** Arthropod Evolution, Arthropod Head Development, Eye Development, Visual System Development

**Email:** [brenda.medina@zoologi.su.se](mailto:brenda.medina@zoologi.su.se)

**Abstract number:** P61

**Title:** Comparative genome annotation for gene family evolution analysis

**Name:** Milena Trabert

**University:** Uppsala University

**Abstract:**

Gene duplications and deletions are common in many eucaryotes, and their effect on phenotypic evolution is of great interest. The statistical framework for these analyses has been developed when the first whole genome sequences became available, and corresponding software is well established. Thanks to the increasing amount of whole genome sequences and annotations available across the whole tree of life, this type of analysis is becoming very accessible and shines new light on many previously unknown examples of phenotypic evolution through gene family expansion or contraction. Unfortunately, the results can be biased through the input data, since it is difficult to distinguish for example if a gene was deleted or merely not annotated in a species. In my work, I am evaluating different annotation methods and different types of reference data for a uniform annotation pipeline to assess the effect of the input data on gene family birth-death dynamics.

**Authors:** Trabert Milena, Uppsala University, Sweden

**Key words:** genome annotation, comparative genomics, gene family evolution

**Email:** milena.trabert@ebc.uu.se

**Abstract number:** P62

**Title:** Visual opsin expression in the damselfly *Ischnura elegans*

**Name:** Natalie Roberts

**University:** Lund University

**Abstract:**

Visual opsins affect color vision in animals. The damselfly, *Ischnura elegans*, relies on vision for mate detection and their eyes are characterized by numerous opsin gene duplicates. Female *I. elegans* express three genetically determined color morphs, with the two most common morphs in Sweden being characterized by blue and green thorax coloration (androchromes and infuscans, respectively). Previous studies have found that exposure to androchrome or infuscans females increases male preference for those morphs and that, in the field, males prefer the most common morph in their population. To assess whether opsin genes are differentially expressed based on previous experience, which may explain male preference based on previous exposure, we quantified opsin gene expression in males that had encountered either androchrome or infuscans females for a 2-day period. Results can explain how visual based mate preferences in this dynamic, polymorphic species.

**Authors:** Roberts, Natalie; Lund University; Sweden Svensson, Erik; Lund University; Sweden

**Key words:** opsin genes, mate choice

**Email:** natalie.roberts@biol.lu.se

**Abstract number:** P63

**Title:** Impact of Domestication, Recent Selection, and Interactions on Behavioural Synchronisation in Dogs and Wolves

**Name:** Lina Roth

**University:** Linköping University

**Abstract:**

This study explored how dogs synchronise their behaviour with their owners, focusing on immediate interactions, breed, and domestication effects. Conducted in outdoor enclosures, the test involved 30 seconds of walking and 30 seconds of standing still. Three studies were performed: Study A tested 20 companion dogs after Ignore, Pet, and Play treatments, showing dogs adjusted their movements to match their owners regardless of treatment. Study B involved 24 ancient breed dogs, 17 solitary hunting dogs, and 20 companion dogs, all synchronising their movements with their owners. Study C tested six socialised pack-living wolves and six pack-living dogs, finding no movement synchronisation with handlers. The results suggest that dogs living with humans synchronise their movements with their owners, while pack-living wolves and dogs do not, indicating that interspecies behavioural synchronisation may depend on the extent of human interaction.

**Authors:** Jasmine Heurlin, György Barabás, Lina S V Roth

**Key words:** Domestication, Behavioural synchronisation, Wolves, Dogs

**Email:** [lina.roth@liu.se](mailto:lina.roth@liu.se)



**Abstract number:** P64

**Title:** New genomic data of the Levant confirm mobility patterns beyond the regions of West Asia

**Name:** Tina Saupe

**University:** Uppsala University

**Abstract:**

With the growing availability of ancient human genomes worldwide, researchers of different disciplines have been increasingly interested in linking changes in genetic-related ancestries, socio-cultural-related patterns, and linguistics focusing particularly on West Asia as the bridge between the geographical regions. Genetic studies have shown that genetic-related ancestries in West Asia were introduced mostly during the Bronze-Iron Age transition, hence the variety of genetic-related ancestries for the North of the Levant is still unclear. Here, we want to present newly generated genetic data of archaeological sites located in the Levant and place them in context with published ancient genomes focusing on possible mobility patterns throughout West Asia. Due to the rather bad preservation of human skeletal remains from this geographical region, we focused firstly on the improvement of the laboratory workflow by applying a target enrichment via in-solution hybridisation on selected individuals. Our first observations of the genetic results have shown a similar picture of genetic-related ancestries supporting previous findings.

**Authors:** Tina Saupe (Human Evolution, IOB, EBC, Uppsala University, Uppsala, Sweden), Carolina Bernhardsson (Human Evolution, IOB, EBC, Uppsala University, Uppsala, Sweden), Mattias Jakobsson (Human Evolution, IOB, EBC, Uppsala University, Uppsala, Sweden)

**Key words:** ancient DNA, Levant, Bronze Age, Target Enrichment

**Email:** tsaupe@live.de

**Abstract number:** P65

**Title:** Heritability of shield size and shape in the green tortoise beetles *Cassida viridis*

**Name:** Erica Winslott

**University:** Lund university

**Abstract:**

The amount of additive genetic variance affects how fast a population can respond to selection, but how additive genetic variance varies over time and space in wild populations is not well understood. One way to assess additive genetic variance is by estimating heritability. We used the green tortoise beetle, *Cassida viridis*, a novel system for studying evolutionary potential in the wild, to address trait and population differences in heritability. Our traits in focus were the size and shape of the extended elytra and pronotum (the shield), which the beetles use as protection from predators. Preliminary results indicate a higher heritability of the beetle shield area than the shield shape. Furthermore, we found potential differences in heritability between two populations, and in some cases these differences depended on the parents' sex. These results suggest that traits and populations might differ in their evolutionary potential.

**Authors:** Winslott, Erica, Lund university, Sweden.

Karlsson Green Kristina, Swedish University of Agricultural Sciences, Sweden.

Lürig Moritz, University of Florida, USA.

Tarka Maja, Lund university, Sweden.

**Key words:** Heritability, Quantitative genetics

**Email:** [erica.winslott@biol.lu.se](mailto:erica.winslott@biol.lu.se)